

FORM PCT/1992
(REV 10-94)

U.S. DEPARTMENT OF COMMERCE PATENT AND TRADEMARK OFFICE

ATTORNEY'S DOCKET NUMBER

TRANSMITTAL LETTER TO THE UNITED STATES
DESIGNATED/ELECTED OFFICE (DO/EO/US)
CONCERNING A FILING UNDER 35 U.S.C. 371

9555.109USWO

U.S. APPLICATION NUMBER (35 U.S.C. 371(c))

Unknown

09/744129

INTERNATIONAL APPLICATION NO.

INTERNATIONAL FILING DATE

PRIORITY DATE CLAIMED

PCT/CA99/00666

21 July 1999

21 July 1998

TITLE OF INVENTION

COMPOSITION FOR ACCELERATING SEED GERMINATION AND PLANT GROWTH

APPLICANT(S) FOR DO/EO/US

SMITH et al.

Applicant herewith submits to the United States Designated/Elected Office (DO/EO/US) the following items and other information:

1. ☒ This is a **FIRST** submission of items concerning a filing under 35 U.S.C. 371.
2. ☐ This is a **SECOND** or **SUBSEQUENT** submission of items concerning a filing under 35 U.S.C. 371.
3. ☒ This express request to begin national examination procedures (35 U.S.C. 371(f)) at any time rather than delay examination until the expiration of the applicable time limit set in 35 U.S.C. 371(b) and PCT Articles 22 and 39(1).
4. ☒ A proper Demand for International Preliminary Examination was made by the 19th month from the earliest claimed priority date.
5. ☒ A copy of the International Application as filed (35 U.S.C. 371(c)(2))
 - a. ☐ is transmitted herewith (required only if not transmitted by the International Bureau).
 - b. ☒ has been transmitted by the International Bureau.
 - c. ☐ is not required, as the application was filed in the United States Receiving Office (RO/US)
6. ☐ A translation of the International Application into English (35 U.S.C. 371(c)(2)).
7. ☒ Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. 371(c)(3))
 - a. ☐ are transmitted herewith (required only if not transmitted by the International Bureau).
 - b. ☒ have been transmitted by the International Bureau.
 - c. ☐ have not been made; however, the time limit for making such amendments has NOT expired.
 - d. ☐ have not been made and will not be made.
8. ☐ A translation of the amendments to the claims under PCT Article 19 (35 U.S.C. 371(c)(3)).
9. ☒ An unsigned oath or declaration of the inventor(s) (35 U.S.C. 371(c)(4)).
10. ☐ A translation of the annexes to the International Preliminary Examination Report under PCT Article 36 (35 U.S.C. 371(c)(5)).

Items 11. to 16. below concern document(s) or information included:

11. ☐ An Information Disclosure Statement under 37 CFR 1.97 and 1.98.
12. ☐ An assignment document for recording. A separate cover sheet in compliance with 37 CFR 3.28 and 3.31 is included.
13. ☒ A FIRST preliminary amendment.
☐ A SECOND of SUBSEQUENT preliminary amendment.
14. ☐ A substitute specification.
15. ☐ A change of power of attorney and/or address letter.
16. ☒ Other items or information: Courtesy copy of PCT/CA99/00666; 1 sheet formal drawing; Form PCT/IB/308; Form PCT/RO/101; Form PCT/ISA/220; Form PCT/ISA/210; Form PCT/IPEA/401; Form PCT/IPEA/408; Form PCT/IPEA/416; Form PCT/IPEA/409

U.S. APPLICATION NO (if known, see 37 CFR 1.5)

Unknown

INTERNATIONAL APPLICATION NO

PCT/CA99/00666

1992 PCT/PTO 7 9 JAN 2001

5555.109USWO

17. [X] The following fees are submitted:

BASIC NATIONAL FEE (37 CFR 1.492(a) (1)-(5)):

Search Report has been prepared by the EPO or JPO.....\$860.00

International preliminary examination fee paid to USPTO

(37 CFR 1.492(a)(1)).....\$690.00

No international preliminary examination fee paid to USPTO (37 CFR 1.482)

but international search fee paid to USPTO (37 CFR 1.445(a)(2)).....\$710.00

Neither international preliminary examination fee (37 CFR 1.482) nor

international search fee (37 CFR 1.445(a)(3)) paid to USPTO.....\$1000.00

International preliminary examination fee paid to USPTO (37 CFR 1.482)

and all claims satisfied provisions of PCT Article 33(2)-(4).....\$100.00

ENTER APPROPRIATE BASIC FEE AMOUNT = \$860.00Surcharge of **\$130.00** for furnishing the oath or declaration later than [] 20 [] 30 months from the earliest claimed priority date (37 CFR 1.492(e)).

\$

CLAIMS	NUMBER FILED	NUMBER EXTRA	RATE	
Total claims	30	-20 = 10	X \$18.00	\$180.00
Independent claims	3	-3 = 0	X \$78.00	\$0.00
MULTIPLE DEPENDENT CLAIM(S) (if applicable)			+ \$260.00	\$

TOTAL OF ABOVE CALCULATIONS = \$1040.00

Reduction by 1/2 for filing by small entity, if applicable. Small entity status is claimed pursuant to 37 CFR 1.27

\$520.00

SUBTOTAL = \$520.00Processing fee of **\$130.00** for furnishing the English translation later than [] 20 [] 30 months from the earliest claimed priority date (37 CFR 1.492(f)).

+

\$

TOTAL NATIONAL FEE = \$520.00

Fee for recording the enclosed assignment (37 CFR 1.21(h)). The assignment must be accompanied by an appropriate cover sheet (37 CFR 3.28, 3.31). \$40.00 per property

+

\$

TOTAL FEES ENCLOSED = \$520.00Amount to be:
refunded \$

charged \$

a. [X] Check(s) in the amount of \$520.00 to cover the above fees is enclosed.b. [] Please charge my Deposit Account No. _____ in the amount of \$ _____ to cover the above fees
A duplicate copy of this sheet is enclosed.c. [X] The Commissioner is hereby authorized to charge any additional fees which may be required, or credit any overpayment to Deposit Account No. 13-2725.**NOTE: Where an appropriate time limit under 37 CFR 1.494 or 1.495 has not been met, a petition to revive (37 CFR 1.137(a) or (b)) must be filed and granted to restore the application to pending status.**

SEND ALL CORRESPONDENCE TO

Gregory A. Sebald

MERCHANT & GOULD

P.O. Box 2903

Minneapolis, MN 55402-0903

SIGNATURE: 

NAME: Gregory A. Sebald

REGISTRATION NUMBER: 33,280

09/744129

JCO2 Rec'd PCT/PTO 1 9 JAN 2001
PATENT

S/N unknown

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Applicant:	SMITH et al.	Docket No.:	9555.109USWO
Serial No.:	unknown	Filed:	concurrent herewith
Int'l Appln No.:	PCTCA0000666	Int'l Filing Date:	21 July 1999
Title:	COMPOSITION FOR ACCELERATING SEED GERMINATION AND PLANT GROWTH		

CERTIFICATE UNDER 37 CFR 1.10

'Express Mail' mailing label number: EL415941482US

Date of Deposit: January 19, 2001

I hereby certify that this correspondence is being deposited with the United States Postal Service 'Express Mail Post Office To Addressee' service under 37 CFR 1.10 on the date indicated above and is addressed to the Assistant Commissioner for Patents, Washington, D.C. 20231

By: 

Name: Grant Miles

PRELIMINARY AMENDMENT

Box PCT
Assistant Commissioner for Patents
Washington, D. C. 20231

Dear Sir:

In connection with the above-identified application filed herewith, please enter the following preliminary amendment in accordance with 37CFR 1.121, a copy of which is enclosed herewith:

IN THE ABSTRACT

Insert the attached Abstract page into the application as the last page thereof.

IN THE SPECIFICATION

A courtesy copy of the present specification is enclosed herewith. However, the World Intellectual Property Office (WIPO) copy should be relied upon if it is already in the U.S. Patent Office.

IN THE CLAIMS

Please cancel claims 22-25 and 33-34 without prejudice or disclaimer.

Please add the following new claims:

--35. (New) The method of claim 17, wherein said composition comprises a bacterial strain which expresses said LCO.

36. (New) The method of claim 35, wherein said bacterial strain is a rhizobial strain.--

Please amend the claims as shown in the attached consolidated claim set.

REMARKS

A new abstract page is supplied to conform to that appearing on the publication page of the WIPO application, but the new Abstract is typed on a separate page as required by U.S. practice.

Applicants respectfully request that the preliminary amendment described herein be entered into the record prior to calculation of the filing fee and prior to examination and consideration of the above-identified application.

Claims 1-21, 26-32 and 35-36 are now pending in the case.

In essence, use claims 1-21 have been replaced by method claims. Further, the claims with multiple-dependencies have been replaced by singly-depending claims. Support for these amendments can be found in the claims as initially filed, as well as throughout the disclosure. Further support can be found, for example, at page 6, lines 4-15; page 8, lines 3-14; page 13, line 27 to page 14, line 3; and Examples 9 and 10. Further support for new claims 35 and 36 can be found, for example, at page 10, lines 23-25; page 12, line 27 to page 13, line 2; and page 8, lines 1-3.


If a telephone conference would be helpful in resolving any issues concerning this communication, please contact Applicants' primary attorney-of record, Gregory A. Sebald (Reg. No. 33,280), at 612.336.4728.

Respectfully submitted,

MERCHANT & GOULD P.C.
P.O. Box 2903
Minneapolis, Minnesota 55402-0903
(612) 332-5300

Dated: January 19, 2001

By



Gregory A. Sebald
Reg. No. 33,280

GAS/vh

ABSTRACT

Lipo Chitooligosaccharide (LCO) [NodBj-V(C18:1, Mefuc)] isolated from *Bradyrhizobium japonicum* strain 532C was able to stimulate seed germination/seedling emergence, or in the case of potato, sprouting, of a number of crop plants representing eight distantly related plant families (Poaceae, Fabaceae, Brassicaceae, Cucurbitaceae, Malvaceae, Asteraceae, Chenopodiaceae and Solanaceae) of plants at 25 and/or at 15°C.

It also promoted sprouting potato minitubers. Other LCOs [NodRM-V(C_{16:2,5}) and LCO from *R. leguminosarum*] were also shown to also display growth-promoting effects on the tested crop plants. The compositions comprising at least one LCO are shown to be effective in promoting growth under both laboratory and field conditions. The invention thus also relates to methods for promoting seed germination and/or seedling emergence and/or growth of plants comprising subjecting the seeds and/or plants to an effective amount of an agricultural composition comprising at least one LCO.

CONSOLIDATED AMENDED CLAIMS

1. Method for enhancing plant crop seed germination and/or seedling emergence and/or growth of a plant crop comprising a treatment in the vicinity of one of a seed, root or plant with a composition which comprises an effective amount of at least one lipo chitoooligosaccharide (LCO) together with an agriculturally suitable carrier, wherein said effective amount enhances seed germination and/or seedling emergence and/or growth of said plant in comparison to an untreated plant.

2. The method according to claim 1, wherein said plant crop is a non-legume.

3. The method according to claim 2, wherein said plant crop is selected from the group consisting of Poaceae, Cucurbitaceae, Malvaceae, Asteraceae, Chenopodiaceae, Solanaceae and Brassicaceae.

4. The method according to claim 3, wherein said plant crop is selected from the group consisting of corn, cotton, cucumber, cantaloupe, lettuce, beet, canola and potato.

5. The method according to claim 1, wherein said LCO is obtainable from a rhizobia selected from the group consisting of *Bradyrhizobium japonicum*, *Rhizobium meliloti* and *Rhizobium leguminosarum*.

6. The method according to claim 5, wherein said LCO is present in said composition at a concentration of between about 10^{-5} M to about 10^{-14} M.

7. The method according to claim 5, wherein said LCO is present in said composition at a concentration of between about 10^{-6} M to about 10^{-12} M.

8. The method according to claim 5, wherein said LCO is present in said composition at a concentration of between about 10^{-7} M to about 10^{-10} M.

9. The method according to claim 1, wherein said composition is effective in enhancing seed germination and/or seedling emergence and/or growth of a plant crop grown under field conditions.

10. The method according to claim 1, wherein said plant is a member of the Fabaceae family.

11. The method according to claim 10, wherein said plant is selected from the group consisting of soybean, bean, alfalfa and clover.

12. The method according to claim 10 [or 11], wherein said LCO is obtainable from a rhizobia selected from the group consisting of *Bradyrhizobium japonicum*, *Rhizobium meliloti* and *Rhizobium leguminosarum*.

13. The method according to claim 12, wherein said LCO is present in said composition at a concentration of between about 10^{-5} M to about 10^{-14} M.

14. The method according to claim 12, wherein said LCO is present in said composition at a concentration of between about 10^{-6} M to about 10^{-12} M.

15. The method according to claim 12, wherein said LCO is present in said composition at a concentration of between about 10^{-7} M to about 10^{-10} M.

16. The method according to claim 10, wherein said composition is effective in enhancing seed germination and/or seedling emergence and/or growth of a plant crop grown under field conditions.

17. A method for breaking the dormancy and/or quiescence of a plant comprising a treatment in the vicinity of a seed, tuber, or root of said plant with an effective amount of an agricultural composition comprising at least one lipo chitooligosaccharide (LCO) together with an agriculturally suitable carrier, wherein said effective amount enables a breaking of the dormancy and/or quiescence of said seed, tuber, or root, in comparison to an untreated seed, tuber, or root.

18. The method according to claim 17, wherein said plant is a member of the family of Solonaceae.

19. The method according to claim 18, wherein said plant is a potato.

20. The method according to claim 19, wherein said growth-promoting activity of said composition enables an increase in yield.

21. The method according to claim 19, wherein said composition further comprises gibberellic acid.

26. A method for enhancing seed germination and/or seedling emergence and/or growth of a plant crop comprising an incubation of a rhizobial strain which expresses a lipo chitooligosaccharide (LCO) in the vicinity of one of a seed and/or root of said plant such that said LCO enhances seed germination and/or seedling emergence and/or growth of said plant crop, wherein said inoculation enhances seed germination

and/or seedling emergence and/or growth in comparison to a non-inoculated seed and/or root of said plant.

27. The method of claim 26, wherein said plant crop is a non-legume.

28. The method of claim 27, wherein said plant crop is selected from the group consisting of Poaceae, Cucurbitaceae, Malvaceae, Asteraceae, Chenopodiaceae, Solanaceae and Brassicaceae.

29. The method of claim 28, wherein said plant crop is selected from the group consisting of corn, cotton, cucumber, cantaloupe, lettuce, beet, canola and potato.


30. The method of claim 26, wherein said rhizobia is selected from *Bradyrhizobium japonicum*, *Rhizobium meliloti* and *Rhizobium leguminosarum*.

31. The method of claim 26, wherein said LCO enhances seed germination and/or seedling emergence and/or growth of said plant grown under field conditions.

32. The method of claim 26, wherein said plant crop is a legume in the Fabaceae family and wherein said LCO enhances seed germination and/or seedling emergence and/or growth of said legume grown under field conditions.

35. The method of claim 17, wherein said composition comprises a bacterial strain which expresses said LCO.

36. The method of claim 35, wherein said bacterial strain is a rhizobial strain.



Xperts
1**TITLE OF THE INVENTION**COMPOSITION FOR ACCELERATING SEED GERMINATION AND PLANT GROWTH

5

FIELD OF THE INVENTION

The present invention relates to agriculture. More specifically, the invention relates to plant seed germination, seedling emergence, quiescence-breakage and plant growth. Even more specifically, the present invention relates to compositions which accelerate plant seed germination, seedling emergence and plant growth of numerous types of crops and to methods using same.

BACKGROUND OF THE INVENTION

Symbiotic microorganisms can promote the growth of legumes by way of biological fixation of nitrogen. More specifically, rhizobiaceae are gram-negative soil bacteria which fix nitrogen and are involved in symbiotic association with these legumes. This symbiotic association between the bacteria and the legume enables the latter to grow in soils having low assimilable nitrogen levels. In return, through photosynthesis, the legume provides the bacteria with the energy it requires to reduce the atmospheric nitrogen into ammonia. This ammonia can then be used by the legume and enters into the nitrogen metabolism. The legume, of the Fabaceae family, forms nodules in which the rhizobia proliferate. The Rhizobiaceae family is in a state of taxonomic flux. It has been reported to comprise four main genera: Rhizobium, Bradyrhizobium, Sinorhizobium and Azorhizobium (U.S. 5,549,718). The symbiotic relationship between nitrogen-fixing bacteria or rhizobia and plants of the Fabaceae family enables the growth of the latter in soils having low levels of available nitrogen, thus reducing the need for nitrogen fertilizers. Since nitrogen fertilizers can significantly increase the cost of crops, and are

associated with a number of polluting effects, biological means to stimulate this symbiotic relationship and/or to decrease the use of nitrogen fertilizers is of great importance.

Initial recognition between *B. japonicum* and soybean involves exchange of molecular signals (Stacey et al, 1995). Legume roots secrete phenolic compounds (Dakora & Philips, 1996; Peters & Verma, 1990), largely from the area of root hair emergence, which act as chemo-attractants to (brady)rhizobia (Nap & Bisseling, 1990), and activate the *nod* genes. Flavones, isoflavones, and flavanones have been identified as the inducing molecules for (brady)rhizobial chemotaxis and for expression of *nod* genes, e.g. genistein, daidzein and several related compounds in soybean (Peters & Verma, 1990). These plant-to-bacteria signal compounds cause expression of the bacterial *nod* (also *nol* and *noe*) genes very rapidly (only a few minutes after exposure) and at very low concentrations (10^{-7} to 10^{-8} M) (Peters et al., 1986). Generally this is through an interaction with *nodD*, which activates the common *nod* genes, although the situation may be more complex, as is the case in *B. japonicum*, where *nodD₁*, *nodD₂* and *nodVW* are involved (Gillette & Elkan 1996; Stacey 1995). *Nod* genes have been identified in the rhizobia that form nitrogen fixing relationships with numbers of the Fabiaceae family (see 5,549,718 and references therein). Recently, the plant-to-bacteria signal molecules have been shown to promote soybean nodulation and nitrogen fixation under cool soil temperatures (CA 2,179,879) and increase the final soybean grain yield on average of 10% in the field and up to 40% under certain conditions.

Among the products of the *nod* genes induced by the plant phenolic signal molecules are various enzymes involved in the synthesis of a series of lipo chitooligosaccharides (LCOs) (Spaink, 1995; Stacey, 1995). These newly synthesized LCOs act as bacterium-to-plant signals, inducing expression of many of the early nodulin genes (Long, 1989). This results in root hair deformation (including curling), cortical cell division leading to initiation of nodule meristems, secretion of additional *nod* gene inducers, and initiation of

infection threads (Verma, 1992). These bacterium-to-plant signals exert a powerful influence over the plant genome and, when added in the absence of the bacteria, can induce the formation of root nodules (Truchet *et al.*, 1991). Thus, the bacteria-to-plant signals can, without the bacteria, induce all the gene activity for nodule organogenesis (Denarie *et al.*, 1996; Heidstra & Bisseling, 1996). Moreover, the above-mentioned activities induced by LCOs can be produced by concentrations as low as 10^{-14} M (Stokkermans *et al.* 1995). The mutual exchange of signals between the bacteria and the plant are essential for the symbiotic interaction. Rhizobia mutants unable to synthesize LCOs will not form nodules. Analysis of the *B. japonicum* *nod* genes indicates that ability to induce soybean nodulation requires at least: 1) a basic tetrameric Nod factor requiring only *nodABC* genes or 2) a pentameric LCO (C18:1, C16:0 or C16: fatty acid and a methyl-fucose at the reducing end, sometimes acetylated) requiring *nodABCZ* genes (Stokkermans *et al.* 1995).

When added to the appropriate legume, LCOs can cause the induction of nodule meristems (Denarie *et al.*, 1996), and therefore cell division activity. One previous publication has shown that LCOs can induce cell cycle activities in a carrot embryogenesis system at levels as low as 10^{-14} M (De Jong *et al.* 1993).

A chemical structure of lipo chitoooligosaccharides, also termed "symbiotic Nod signals" or "Nod factor", has been described in U.S. Patents 5,549,718 and 5,175,149. These Nod factors have the properties of a lectin ligand or lipo-oligosaccharide substances which can be purified from bacteria or synthesized or produced by genetic engineering.

The relationship between environmental variables, such as low root zone temperature (RZT) and pH, and the interplay of molecular signals has only recently become a subject of investigation. For example, some soybean genotypes have less synthesis abilities for isoflavones under cool soil temperature, whereas a higher isoflavone concentration is needed to turn on the *nod* genes of *B. japonicum* (Zhang and Smith 1995 and 1997). The plant-to-

bacteria signal molecules (i.e. isoflavones) have been shown, among other things, to overcome the negative effect of low temperature on the early events of symbiotic nitrogen fixation (Canadian application number 2,179,879).

While the effects of plant-to-bacteria signal molecules (i.e. isoflavones) on nodulation, nitrogen fixation, growth and protein yield of legumes, such as soybean, and on bacteria-to-plant signal molecules (LCOs) on nodulation and nitrogen fixation in legumes have been described under certain conditions, the effect of the bacteria-to-plant signal molecules on the growth of non-legumes is unknown. In fact, the role of such bacteria-to-plant signal molecules on non-legumes has never been assessed. In addition, the effect of LCOs on processes other than nodulation of legumes has yet to be studied.

There thus remains a need to assess the effect of LCOs on seed germination, seedling emergence and/or growth of plants in general and especially of non-legume plants.

The present invention seeks to meet these and other needs.

The present description refers to a number of documents, the contents of which are herein incorporated by reference in their entirety.

SUMMARY OF THE INVENTION

The invention concerns a composition for enhancing seed germination, seedling emergence and growth of plants and especially of crop plants. More specifically, the present invention relates to a composition comprising an LCO which can increase seed germination and/or seedling emergence and/or growth of a legume, in addition to acting as a trigger to initiate legume symbiotic nitrogen fixation. More particularly, the invention relates to increased seed germination and/or seedling emergence and/or growth of soybean, pea and red clover.

Surprisingly, the compositions of the present invention act not only on legumes such as soybean but on plants in general as exemplified with

non-legume crops from different plant families Poaceae, Cucurbitaceae, Malvaceae, Asteraceae, Chenopodiaceae and Solonaceae. More specifically, the non-legume crops exemplified herein include corn, cotton, cantaloupe, cucumber, canola, lettuce, potato and beet. The present invention thus also
5 refers to compositions for enhancing seed germination and/or seedling emergence and/or growth of non-legumes. More particularly, the invention relates to compositions comprising an LCO for enhancing seed germination, seedling emergence and growth of non-legumes. Non-limiting examples of such non-legumes include cotton, corn, canola, potato, cucumber, cantaloupe, lettuce
10 and beet. Broadly, the present invention relates to compositions comprising an LCO for promoting growth of a crop. Non-limiting examples of crop plants include monocot, dicot, members of the grass family (containing the cereals), and legumes.

Thus, the present invention relates to agricultural
15 compositions comprising at least one LCO (and methods of using same) for promoting seed germination, and/or early development of seedlings, and/or emergence of sprouts from tubers, and/or rapid development of new plants from higher plant perinating structures.

In a particular set of experiments in the field, a composition
20 of the present invention comprising an LCO was shown to significantly enhance early plant growth.

The invention in addition relates to methods for enhancing seed germination and/or seedling emergence and/or growth of plants and/or for breaking the dormancy thereof comprising a treatment in the vicinity of a seed
25 or seedling or plant with an effective amount of an agricultural composition comprising an LCO and an agriculturally suitable carrier for a sufficient time and under conditions which enable an increased germination of the seed and/or an increased emergence of the seedling and/or an increased growth of the plant and/or a triggering of the growth of a dormant plant.

The invention also relates to compositions and methods for breaking the dormancy of a plant and initiating the growth thereof. In a particular embodiment, the invention relates to the breaking of dormancy of potato.

- 5 The Applicant is the first to show that a composition comprising an LCO can have a significant effect on seed germination, and/or seedling emergence of legumes. Moreover, the Applicant is the first to show the surprising effect of signal molecules involved in bacteria-legume signalling on the growth of non-legume plants. In addition, the Applicant is the first to show
- 10 that a composition comprising an LCO had an effect on non-legume seed germination and/or seedling emergence and/or plant growth of the non-legume. Also, the Applicant is the first to show that an LCO can not only act as a dormancy breaker but that it can also significantly increase the yield of a dormant plant following the dormancy breakage, when compared to known
- 15 dormancy breakers.

- While the seed germination and/or seedling emergence and/or plant growth enhancing capabilities of the compositions of the instant invention are demonstrated with corn, cotton, canola, potato, cantaloupe, lettuce, beet, cucumber, soybean, pea and red clover, they are applicable to
- 20 plants in general and more especially to crop plants. Indeed, the plants chosen for the experiments presented herein are crops from significantly divergent plants in eight distinct families: (1) corn, the only monocot tested herein, in the family of grasses (Poaceae), which also contains the cereals; (2) cucumber and cantaloupe, the latter being a plant used horticulturally, and being slow to
- 25 germinate at low temperature [its base temperature is about 14°C] (Cucurbitaceae); (3) cotton, one of the most important fibre crops on the planet (Malvaceae); (4) lettuce (Asteraceae); (5) beet (Chenopodiaceae); (6) potato, a very important crop (Solonaceae, which also includes tobacco, peppers and tomato); and two families of legumes (7) canola, representing the mustard group
- 30 (Brassicaceae) and (8) soybean (representative of oil seed crop), bean


(representative of a crop for human consumption) and red clover and alfalfa (forage legumes) (all of the Fabaceae family).

In view of the diversity of the plants tested, and of the similar results obtained with these different crop plants, it can be predicted that such results will apply to crop plants in general. It follows that a person skilled in the art can adapt the teachings of the present invention to other crops. Non-limiting examples thereof include tobacco, tomato, wheat, barley, rice, sunflower and plants grown for flower production (daisy, carnation, pansy, gladiola, lilies and the like). It will be understood that the compositions can be adapted to specific crops, to meet particular needs.

In accordance with the present invention, there is thus provided an agricultural composition for enhancing plant crop seed germination and/or seedling emergence and/or growth of a plant crop comprising a growth-promoting amount of at least one lipo chitooligosaccharide (LCO) together with an agriculturally suitable carrier. There is also provided a composition for breaking the dormancy and/or quiescence of a plant, comprising a growth-promoting amount of at least one lipo chitooligosaccharide (LCO) together with an agriculturally suitable carrier. Furthermore, there is provided a method for enhancing seed germination and/or seedling emergence and/or growth of a plant, comprising a treatment in the vicinity of one of a seed, root or plant with a composition comprising an agriculturally effective amount of a lipo chitooligosaccharide (LCO) in admixture with an agriculturally suitable carrier medium, wherein the effective amount enhances seed germination and/or seedling emergence and/or growth of the plant in comparison to an untreated plant. There is further provided a method for enhancing seed germination and/or seedling emergence and/or growth of a plant crop comprising incubating a rhizobial strain which expresses a lipo chitooligosaccharide (LCO) in the vicinity of one of a seed and/or root of the plant such that the LCO enhances seed germination and/or seedling emergence and/or growth of the plant crop.

As used herein, the term "rhizobia" is used broadly to refer to bacterial strains which are involved in a nitrogen fixing symbiotic relationship with a legume.

As used herein, the term "LCO" refers broadly to a Nod factor which is under the control of at least one nodulation gene (*nod* gene), common to rhizobia. LCO therefore relates to a bacteria-to-plant signal molecule which induces the formation of nodules in legumes and enables the symbiotic bacteria to colonize same. Broadly, LCOs are lipo chitooligosaccharide signal molecules, acting as phytohormones, comprising an oligosaccharide moiety having a fatty acid condensed at one of its end. An example of an LCO is presented below as formula 1




in which:

G is a hexosamine which can be substituted, for example, by an acetyl group on the nitrogen, a sulfate group, an acetyl group and/or an ether group on an oxygen,

R_1 , R_2 , R_3 , R_5 , R_6 and R_7 , which may be identical or different, represent H, $\text{CH}_3\text{CO}-$, $\text{C}_x\text{H}_y\text{CO}-$ where x is an integer between 0 and 17, and y is an integer between 1 and 35, or any other acyl group such as for example a carbamyl,

More specific LCOs from *R. meliloti* have also been described
5 in 5,549,718 as having the formula II



25 Even more specific LCOs include NodRM, NodRM-1, NodRM-
3. When acetylated (the R = $\text{CH}_3\text{CO}-$), they become AcNodRM-1, and
AcNodRM-3, respectively (U.S. 5,545,718).

LCOs from *B. japonicum* have also been characterized in U.S. 5,175,149 and 5,321,011. Broadly, they are pentasaccharide phytohormones comprising methylfucose. A number of these *B. japonicum*-derived LCOs are

described : BjNod-V ($C_{18:1}$); BjNod-V (A_C , $C_{18:1}$), BjNod-V ($C_{16:1}$); and BjNod-V (A_C , $C_{16:0}$), with "V" indicating the presence of five N-acetylglucosamines; "Ac" an acetylation; the number following the "C" indicating the number of carbons in the fatty acid side chain; and the number following the ":" the number of double bonds.

It shall also be understood that compositions comprising different LCOs, are encompassed within the scope of the present invention. Indeed, while the present invention is exemplified with NodBj-V($C_{18:1}$)₁₁ also known as BjNod-V($C_{18:1}$ MeFuc); NodRM-V($C_{16:2}$, S); and NodRI, any LCO produced by a rhizobia which is capable of entering into a nitrogen fixation relationship with a legume (i.e. a member of the Fabiaceae family) is expected to have the potential to show the same properties as those described herein. It will be clear to the person of ordinary skill that the selection of a rhizobia known to be expressing LCOs at high levels, or known to express an LCO having an effect on a broader spectrum of legumes could be advantageous.

It will also be clear that the LCO compositions of the present invention could also comprise more than one signal molecule. Non-limiting examples of such compositions include agricultural compositions comprising in addition to one LCO: (1) at least one additional LCO; (2) at least one plant-to-bacteria signal molecule; (3) gibberellic acid or other agents or compounds known to promote growth or fitness of plants; and mixtures of such compositions (1), (2) or (3).

It shall be clear that having identified new uses for LCO, bacteria could be genetically engineered to express *nod* genes and used for producing LCOs or for direct administration to the plants and/or seeds.

Thus, while the instant invention is demonstrated in particular with LCOs from *Bradyrhizobium japonicum*, *Rhizobium meliloti* and *R. leguminosarum* and selected legumes and non-legume crops, the invention is not so limited. Other legume crops, non-legume crops and rhizobial strains may

be used using the same principles taught herein. Preferred matching of rhizobia with legume crop groups include, for example:

<u>rhizobial species</u>	<u>Legume crop group</u>
<i>R. meliloti</i>	alfalfa, sweet clover
<i>R. leguminosarum</i>	peas, lentils
<i>R. phaeosolii</i>	beans
<i>Bradyrhizobium japonicum</i>	soybeans
<i>R. trifolii</i>	red clover

As will be apparent to the person of ordinary skill to which the present invention is directed, the growth-stimulating compositions of the present invention can be applied to other crop plants and especially to other warm climate adapted crop plants (plants or crops having evolved under warm conditions [i.e. tropical, subtropical or warm temperature zones] and whose metabolism is optimized for such climates). It should be understood that the growth-enhancing compositions of the present invention should find utility whenever a particular crop is grown in a condition which limits its growth. More particularly, whenever a particular plant crop is grown at a temperature which is below its optimum temperature for seed germination, seedling emergence, growth and the like. Such temperatures are known in the art. For example, optimum temperatures for germination of corn, soybean, rice and cotton are 30°C, 34-36°C, 30-32°C, and 34°C, respectively. The minimum germination temperatures (or base temperatures) for these crops are 9°C, 4°C, 8 to 10°C, and 14°C, respectively, while the maximum germination temperatures are 40°C, 42-44°C, 44°C and 37°C, respectively. The compositions of the present invention therefore find utility, among other things, in enhancing germination of warm climate adapted crops when grown at temperatures between their base temperature for seed germination, and/or seedling emergence and/or growth and their optimum temperature for germination. The compositions of the present invention find utility in general in enhancing seed germination and/or seedling emergence and/or growth of crop plants when grown under conditions which

delay or inhibit seed germination and/or seedling emergence thereof. Non-limiting examples of such inhibiting conditions (as known from their signalling inhibition in bacteria-legume interactions, their inhibition or delay of the bacteria-plant symbiotic relationship) include pH stress, heat-stress, and water stress.

5 It will be nevertheless recognized that the compositions and methods of the present invention enhance growth of plants grown under optimal conditions.

Thus, the compositions and methods of the present invention should not be limited to plants growing under sub-optimal conditions.

10 The term "environmental conditions which inhibit or delay the bacterial-plant symbiotic relationship" should be interpreted herein as designating environmental conditions which postpone or inhibit the production and exchange of signal molecules between same and include, without being limited thereto: conditions that stress the plant, such as temperature stress,
15 water stress, pH stress as well as inhibitory soil nitrogen concentrations or fixed nitrogen.

"An agriculturally effective amount of a composition" for increasing the growth of crop plants in accordance with the present invention refers to a quantity which is sufficient to result in a statistically significant
20 enhancement of growth and/or of protein yield and/or of grain yield of the plant crop as compared to the growth, protein yield and grain yield of the control-treated plant crop. As will be seen below, the growth promoting activity of the LCOs are observable over a broad range of concentrations. Indeed, LCO growth-promoting activities can be observed at an applied concentration of
25 about 10^{-5} to 10^{-14} M, preferably about 10^{-6} to about 10^{-12} M and more preferably about 10^{-7} to about 10^{-10} M.

The term "immediate vicinity of a seed or roots" refers to any location of a seed or roots wherein if any soluble material or composition is so placed, any exhibit of the plant or of the bacteria, or bacterial cells will be in

actual contact with the seed as it germinates or the roots as they grow and develop.

5 Direct or indirect methods of inoculation with the composition of the present invention can be employed. During direct inoculation the composition is applied directly to the seed prior to sowing. This can most simply be accomplished by spraying the seed with or dipping the seed into a liquid culture containing the desired components.

10 The recitation "short season condition" refers herein broadly to temperatures of the middle and temperate zones and shorter. Typically, the active growing season is around 1/2 to 2/3 of the year. Short season conditions broadly refers to a frost-free period of less than half the year, often on the order of 100 frost-free days.

15 By "nodulation gene-inducing" or "*nod* gene-inducing" is meant bacterial genes involved in nodule establishment and function.

By "seed germination" is meant a clear evidence of root growth developing from the embryo on the seed. When referring to an "increased seed germination", the Applicant refers to a significant difference in seed germination between the treated versus the control seed.

20 "Seedling emergence" is meant to refer to growth of the plant which is observable above the rooting medium surface. When referring to an "enhanced seedling emergence", the Applicant refers to a significant observable difference between the growth of the seedling in the treated versus the control.

BRIEF DESCRIPTION OF THE DRAWING

25 Having thus generally described the invention, reference will now be made to the accompanying drawing, showing by way of illustration a preferred embodiment thereof, and in which:

Figure 1 shows the seed germination enhancing effect of a composition according to the present invention on corn.

Other objects, advantages and features of the present invention will become more apparent upon reading of the following non-restrictive description of preferred embodiments with reference to the accompanying drawing which is exemplary and should not be interpreted as limiting the scope of the present invention.

DESCRIPTION OF THE PREFERRED EMBODIMENT

While the effects on nodulation were detected upon treatment of soybean with SoyaSignal® (a composition comprising both the plant-to-bacteria and bacteria-to-plant signal molecules), it was also noted that in many of the field experiments the plants that received some sort of genistein treatment emerged from the soil sooner. Thus, an experiment, in which genistein alone, *B. japonicum* alone, and genistein plus *B. japonicum* were added to soybean seeds under controlled environment conditions, was conducted. Because slow germination of corn (and other plants, as well) is a serious agricultural problem in eastern Canada because the weather conditions limit the growth thereof, corn was also treated in a similar fashion. The experiment showed that the seed germination and seedling emergence promoting effect was present with the combination of genistein plus *B. japonicum*, leading to the conclusion that the enhancing effects were due to the LCOs produced by genistein exposed *B. japonicum*. Purification (HPLC and otherwise) of the LCO most abundantly produced by genistein-stimulated *B. japonicum* (NodBj-V(C_{18:1a}, 11)) was carried out. This was aided by the gracious gift of enough LCO material to standardize the assay (G. Stacey, University of Tennessee at Knoxville; U.S. 5,175,149 and U.S. 5,321,011) which allowed both isolation and quantification. With isolated NodBj-V(C_{18:1a}, 11), research on the ability of this compound to stimulate seed germination, seedling emergence and growth of leguminous and non-leguminous plants could be conducted.

These experiments surprisingly demonstrated that the addition of SoyaSignal® (which comprises both an isoflavone and an LCO; the

latter at a concentration of about 10^{-5} M) accelerates the germination of corn seeds, whereas isoflavone solutions alone do not. Presumably this effect was due to the LCOs produced by *B. japonicum* cells and induced by the presence of isoflavones. When the seedlings were harvested (still at the mesocotyl stage) they were 44% longer and 33% as heavier in the genistein-*B. japonicum* treated versus non-treated plants (Figure 1). In addition, not only did seedling emergence increase, but the rate of cotton seed germination was also accelerated by the application of SoyaSignal®. The germination rate of cotton seeds treated with SoyaSignal® (0.66 ml/kg seed) increased by 145% compared to those control seeds that were treated with pure water. Both the corn and cotton experiments were conducted at low temperatures, 15°C and 17°C for corn seeds and for cotton seeds, respectively.

The field trial showed that the time of tasselling of sweet corn treated with SoyaSignal® (planted on May 6 on the Experimental Farm of McGill University, Quebec) was 1 to 2 days earlier compared to that of untreated plants. Soybean seeds that received SoyaSignal® (planted on June 22 in Martinsville, Illinois) emerged 8 hours earlier compared to control seeds while the first trifoliar fully expanded 1 day earlier. At the agronomy farm of Purdue University, Indiana, soybeans planted in early June and observed in early July were already one stage further in their development (V6) compared to the control plants (V5). In a farmer trial (in Jackson, Illinois), plants that received SoyaSignal® had many more nodules on the secondary roots and were 10% taller than untreated plants.

Thus, an LCO (a bacteria-to-plant signal molecule involved in the establishment of the symbiotic relationship between a rhizobia and a legume) can promote growth of corn, a monocot distantly related to legumes. Based on the evolutionary divergence of corn from legumes and the significant response thereof to the LCO treatments, corn was used as a model plant system in follow-up experiments. These experiments demonstrated that the results obtained with corn were also observable with all other crop plants tested.

Taken together, the laboratory data and field trials presented herein show that an LCO can increase seed germination, seedling emergence and plant growth of legumes and non-legume plants under controlled environment and field conditions.

5 The signal molecules are also shown to break the dormancy of potato tubers. Of note, the dormancy experiments showed that the signal solution was better at increasing the yield of potato tubers as compared to other dormancy breakers (i.e. gibberellic acid).

10 The precise mechanism of action of LCOs on seed germination, seedling emergence, dormancy and plant growth of legumes and non-legumes is not fully understood. The general understanding of the role of LCOs in signalling during the establishment of the legume-rhizobia symbiosis was described above. When added to the appropriate legume, LCOs can cause the induction of nodule meristems. Thus, it is possible that LCOs might be
15 normal signal molecules in higher plants, so that exogenously supplying them simply increases their levels and, therefore, the activity of the things they would normally regulate. Alternatively, there may be an endogenous class of signal molecules which play important roles in plant development, and have a conformation similar to those of LCOs. One possible candidate for this is the oligosaccharins (Fry *et al.* 1993), some of which do stimulate meristem activity (Pavlova *et al.* 1992). LCOs are somewhat similar in structure and chemistry to the oligosaccharins (Fry *et al.* 1993) and can, in the broadest sense, be included in that group (Stokkermans *et al.* 1995). However, the signal molecules with a
20 similar conformation need not be chemically similar, as demonstrated by the ability of opiates (plant alkaloids) to fit into receptor sites normally occupied by endorphins (oligo-peptides). Nothing is known regarding the mechanisms by which LCOs cause this activity. Without being limited to a particular theory, the present invention is nevertheless the first to have identified a seed germination, and/or seedling emergence and/or dormancy breaking and/or plant growth
25 promoting effect of a composition comprising LCOs on non-legume plants.
30

Crops, such as soybean, corn and cotton evolved in relatively warm climates and, as a result, have high base temperatures for germination, being of about 5°C for soybean, 10°C for corn and 14°C for cotton. These high base germination temperatures lead to slow emergence after planting, resulting in slow leaf ground cover early in the season (when the temperature is sub-optimal), which in turn leads to poorer early season light interception, poorer competition with weeds (and therefore greater need for herbicide application) and increased soil erosion during heavy rainfall events. To simplify, these crops are often grown under conditions which limit their seed germination, and/or seedling emergence and/or growth. Hence, the use of a growth-promoting factor which is in limiting amount can compensate for a deficiency or stress in the growth conditions. Using SoyaSignal® as a plant growth regulator could thus partially overcome the negative effects of environmental stress conditions, such as low soil temperature on crop seed germination, seedling emergence and plant development. Thus, the present invention provides the means to improve the production of crops of tropical and subtropical origin in the temperate zones and may extend their production into shorter season areas. In addition, the present invention provides the means to improve production of crops growing under stress conditions.

The present invention is illustrated in further detail by the following non-limiting examples.

EXAMPLE 1

Induction of LCO production by *Bradyrhizobium japonicum*

The first culture containing *Bradyrhizobium japonicum* (strain 532C) was grown at 28°C in 100 -125 mL of sterile yeast mannitol media (YEM) with pH 6.8, shaken at 150 rpm until the OD₆₂₀ reaches 0.4-0.6 (4-6 days). Thereafter, a 2L bacterial subculture was started by inoculating with material from the first culture (5 mL of first culture per 250 mL of YEM media), for 5-7 days (OD₆₂₀ - 0.8-1.0), as above. At this stage, 0.25 L of 50 µM genistein (in

methanol) were added to each 250 mL of bacterial subculture (genistein concentration of 5 μM) and the culture was incubated for 48-96 hours, the flavone thereby inducing LCO production in the bacterial cells.

5

EXAMPLE 2

Induction of LCO production by *Rhizobium meliloti* or *Rhizobium leguminosarum*

10

The first culture of *Rhizobium meliloti* strain RCR2011 was grown at 28°C in 100-125 mL of sterile yeast mannitol media (YEM) with pH 6.8, shaken at 150 rpm until the OD₆₂₀ reaches 0.4-0.6 (2-3 days). Thereafter, a 2L bacterial subculture was started by inoculating first culture (5 mL of first culture per 250 mL of YEM media), for 2-3 days (OD₆₂₀ - 0.8-1.0), as above. At this stage, 0.25 mL of 50 μM luteolin (in methanol) was added to each 250 mL of bacterial subculture (luteolin concentration of 5 μM) and the culture was incubated for 48 hours, the flavone thereby inducing LCO production in the bacterial cells.

15

20

For LCO production by *Rhizobium leguminosarum*, the rhizobia was grown similarly as above. The flavone (naringenin) was added to the subculture of *R. leguminosarum* (10 μM) and the procedure carried out as above.

EXAMPLE 3

Extraction and purification of LCOs

25

Two liters of bacterial subculture were phase-partitioned against 0.8 L of HPLC-grade 1-butanol by shaking overnight. The upper butanol layer was then transferred to a 1 L evaporation flask and evaporated at 80°C to 2-3 mL of light brown, viscose material with a Yamato RE500 Rotary Evaporator, which was resuspended in 4 mL of 18% acetonitrile and kept in the dark at 4 °C in a sealed glass vial.

HPLC analysis was conducted with a Vydac C18 reversed-phase column with flow rate 1.0 mL/min and a Vydac guard column. As a baseline, acetonitrile (AcN/ H₂O; w/w) was run through the system for at least 10 min. When the sample was loaded, an isocratic elution was started by 18% of AcN for 45 min. This step aims at removing all non-polar contaminant light fractions. Thereafter, a gradient elution for 90 min. with 18-82% AcN was performed. LCOs began to elute after 94-96 min. of HPLC run time.

For the purification of LCOs from *R. leguminosarum* (which nodulates numerous legumes), the HPLC peaks were identified and compared to those obtained with *B. japonicum* and *R. meliloti*. LCO peaks which were different from those of these two other rhizobia were identified collected. Thus, it is strongly suggested that the *R. leguminosarum* LCOs used herein are different from that of *B. japonicum* or *R. meliloti*.

EXAMPLE 4

Effect of LCO on emergence of some plant species

Plastic pots (7.5 cm dia) were filled with 15g of autoclaved vermiculite. Seeds of corn (*Zea mays* - Poaceae), bean (*Phaseolus vulgaris* - Fabaceae), canola (*Brassica napus* - Brassicaceae), cucumber (*Cucumis sativus* - Cucurbitaceae), cantaloupe (*Cucumis melo* - Cucurbitaceae), cotton (*Gossypium* sp.- Malvaceae), lettuce (*Lactuca sativa* - Asteraceae), beet (*Beta vulgaris* - Chenopodiaceae), and soybean (*Glycine max* - Fabaceae), were placed at 2.5 cm deep at the rate of 5 or 10 seeds per pot. Pots were irrigated with either 25 mL of LCO solution at different concentrations (10^{-6} – 10^{-14} M) or aqueous acetonitrile or water, as controls. Acetonitrile was included as one of the controls since LCO was purified in this solvent (see Example 3), after 4 days the pots were irrigated with 10 mL of water once every two days. Each treatment had 4 replications in a randomized block design. Pots were placed on a green house bench maintained at $25 \pm 2^{\circ}\text{C}$ with a day/night cycle 16/8h and relative humidity of 70%, or in a growth chamber set at 15°C with a 16:8 day/night cycle.

As defined above, seed germination has occurred when clear evidence of root growth developing from the embryo on the seed is observed.

As the time required for seedling emergence of the species used in the experiment varied considerably, observation on seedling emergence was recorded when the emergence was observed for at least 50% in most of the treatments. As defined above, seedling emergence has occurred when growth of the plant can be observed above the rooting medium surface. The percent emergence was calculated. The data were analyzed with Statistical Analysis System, version 6.12 (SAS institute Inc. Cary, NC, USA).

LCO treatment reduced the time required from sowing to emergence of a number of economically important plant species tested. Among the species tested, *Z. mays*, *L. sativa*, *B. vulgaris*, *P. vulgaris*, and *G. max* showed significant increases in emergence when treated with LCO at 25°C (Table 1), while, *C. sativus* and *B. napus* showed similar effects at 15°C (Table 2).

TABLE 1
Effect of lipo chitooligosaccharide on
seedling emergence (%) at 25°C

Treatment	Zea mays	Beta vulgaris	Glycine max	Gossy- pium sp.	Cucumis melo	Letuca sativa	Phaseolus vulgaris
Control	40 c ^w	33 a	40 e	55 b	80 a	5 d	44 abc
10 ⁻⁶ M	76 ab	NT	65 d	88 a	NT	45 a	67 abc
10 ⁻⁷ M	68 abc	66.6	80 bc	66 ab	100 b	35 ab	89 a
10 ⁻⁸ M	84 a	NT	90 ab	88 a	NT	10 dc	78 ab
10 ⁻⁹ M	88 a	66 b	100 a	88 a	100 b	20 bcd	67 abc
10 ⁻¹⁰ M	84 a	NT	70 cd	88 a	NT	25 abcd	44 abc
10 ⁻¹¹ M	68 abc	86 b	50 e	NT	100 b	26 abcd	22 c
10 ⁻¹² M	48 abc	NT	80 bc	NT	NT	30 abc	33 bc
10 ⁻¹³ M	40 c	80 b	70 cd	NT	100 b	5 d	33 bc
10 ⁻¹⁴ M	40 c	NT	70 cd	NT	NT	10 d	33 bc

^w means with in the same column, followed by the same letter are not significantly different ($p < 0.05$) by ANOVA protected LSD test.

NT- Not tested

TABLE 2

Effect of lipo chitooligosaccharide
on seedling emergence at 15°C

Treatment	<i>Cucumis sativus</i> 60 c [¶]	<i>Brassica napus</i> 32.5 c
Lipo chitooligosaccharide		
10 ⁻⁶ M	65 abc	35 bc
10 ⁻⁷ M	85 ab	32 c
10 ⁻⁸ M	80 ab	35 bc
10 ⁻⁹ M	70 abc	52 ab
10 ⁻¹⁰ M	50 c	62 a
10 ⁻¹¹ M	80 ab	47 bc
10 ⁻¹² M	80 ab	45 bc
10 ⁻¹³ M	70 abc	37 bc
10 ⁻¹⁴ M	70 abc	30 c

¶ means with in the same column, followed by the same letter are not significantly different ($p \leq 0.05$) by ANOVA protected LSD test.

In some plants, germination/emergence promoting effects of LCOs is seen at all temperatures suitable for growth, while in others, it is only observed under temperature-limiting conditions.

5

EXAMPLE 5**Effects of LCO on early growth of corn**

The percent seedling emergence was recorded at 4 days after sowing (DAS). Plant height was recorded from 4 DAS to 15 DAS. Plants were harvested at 15 DAS and leaf area, root length and the number of roots per plant were recorded. The plants were then dissected and placed in paper covers and dried at 90°C for 24h and the dry weights of roots, shoots and the spent seeds recorded. The data were analyzed with the Statistical Analysis System version 6.12 (SAS institute Inc. Cary, NC, USA).

In LCO treatments, seedling emergence started 2-3 days after seeding while in the control it was 3-4 days. LCO treatments significantly increased leaf area, root length, number of roots, shoot dry weight and root dry weight, while the weight of spent seed recorded significant decreases as compared to the control (Table 3). The optimum effect was observed at an LCO concentration of 10^{-6} M. The decrease in spent seed weight is attributed to the rapid translocation of stored reserve from the seed endosperm to the embryo. Of interest, a dramatic increase of α -amylase activity was observed in the treated seeds.

TABLE 3

Effect of lipo chitooligosaccharide on early growth (after seedling emergence) of *Zea mays*

Treatment	Leaf area (cm ²)	Root length (mm)	No. of roots	Plant height at 15 DAS (mm)	Root dry wt./plant (mg)	Spent seed dry wt. (mg)	Shoot dry wt./plant (mg)
Water	5.81 f ^v	103.1 e	5.1 e	79.8d	53.2 de	163.6 a	28.8 e
Acetonitrile	7.63 ef	106.3 e	5.8 de	94.1cd	65.4 e	147.4 ab	30.1 de
LCO 10 ⁻⁵	11.7 cd	137.0 cd	6.8 cd	102.9c	75.7 cd	123.5 bc	41.6 bcd
LCO 10 ⁻⁶	18.2 a	150.8 bc	7.66 abc	130.4ab	85.1 bc	90.7 cd	51.4 b
LCO 10 ⁻⁷	15.7 ab	153.9 b	8.0 ab	130.1ab	96.5 ab	99.5 cd	51.6 b
LCO 10 ⁻⁸	19.3 a	187.0 a	8.4 a	142.6a	103.4 a	76.3 d	66.2 a
LCO 10 ⁻⁹	17.0 ab	144.4 bcd	7.75 abc	134.7ab	83.3 bc	93.7 cd	46.8 bc
LCO 10 ⁻¹⁰	13.9 bc	149.1 bc	7.95 ab	127.1b	89.5 abc	89.8 cd	51.4 b
LCO 10 ⁻¹¹	8.51 def	132.1 d	6.89 cd	103.3	74.6 cd	115.9 bc	38.0 cde
LCO 10 ⁻¹²	8.0 def	137.9 cd	7.25 d	98.3c	84.6 bc	105.4 cd	40.6 bcde
LCO 10 ⁻¹³	10.4 cde	130.9 d	6.7 cd	91.8cd	85.4 bc	109.6 cd	43.6 bc
LCO 10 ⁻¹⁴	11.2 cde	136.8 cd	7.36 abc	99.9c	90.2 abc	102.9 cd	46.6 bc

^v means with in the same column, followed by the same letter are not significantly different ($p \leq 0.05$) by ANOVA protected LSD test.

Taken together, Examples 4 and 5 and Tables 1-3 show that LCOs can stimulate seedling emergence in all tested plants. In addition, a significant growth stimulation of corn was observed. Furthermore, the spent seed weight results suggested that LCOs also had an effect on seed germination for all tested plants. The growth-promoting effect of LCOs on corn, a plant quite distantly related to legumes (i.e. corn is a monocot), strongly suggests that plants in general should show the same growth-responses to LCO treatment.

EXAMPLE 6

10 Dormancy breaking activity of LCO on potato mini tubers

Signal solution is a bacterial fermentation tank product, comprising approximately 10^{-6} M LCO from *B. japonicum*. More specifically, signal solution is the supernatant from a culture of *B. japonicum* in which genestein (a flavone) had been introduced to promote LCO expression. Following the subculture of *B. japonicum*, the bacteria was removed. While the stimulatory effect of Signal solution in the soybean-*Bradyrhizobium japonicum* complex has been described (Zhang and Smith, 1995; Zhang *et al.* 1996), the effects of these plant substances in other plant species and their associated rhizospheres' organisms have not been investigated.

20 Gibberellic acid (GA) and kinetin affect both the germination rate and the percent germination of crop seeds.

Some studies have indicated that plant growth regulators (PGRs), such as gibberellic acids (GAs), stimulate seed germination at low temperatures. Durrant and Mash (1991) reported that adding gibberellins (GA₄₇) to sugar-beet seeds (*Beta vulgaris* L. Var. *altissima*) was beneficial to seed germination under cold, wet conditions.

25 Kepczynski and Bialecka reported that Methyl jasmonate (JA-Me) inhibited or retarded germination of *Amaranthus caudatus* seeds in darkness at 24°C. Ethephon, ACC (1-aminocyclopropane-1-carboxylic acid) and gibberellins

(GA₃ or GA₄₊₇) partially or completely reversed this inhibition depending on the concentration of JA-Me applied.

Indeed, gibberellic acid, as well as bromoethane, are used commercially to break dormancy and to stimulate sprout formation.

- 5 Treatments were carried out on microtubers (200-400 mg) that had been cold-stored for 8 wk to determine their effect on breaking dormancy. Signal solution was used at full strength (100%) or diluted to 20% (as for soybean), 12%, or 6% of full strength. GA₃ (500 mg l⁻¹), water soaking, and control treatments were performed for comparison purposes. Microtuber soaking treatments lasted 24
- 10 h and then incubation occurred either in the light (40 μ mol m⁻²s⁻¹ cool-white fluorescent) or in the dark. Five microtubers were used in each treatment. Observations for sprouted microtubers were made at 1 and 2 wk.

- One hundred % signal solution was as effective as GA₃ (500 mg l⁻¹) when evaluated after 1 wk with respect to the number of sprouted microtubers.
- 15 Table 4 shows the effect of signal solution (SS) on dormancy breaking of potato microtubers as compared to the known dormancy breaker gibberellic acid (GA₃) (200-450 mg) that had been cold-stored at 5°C for 8 weeks and evaluated after treatment and incubation with or without light for 1 and 2 weeks, for number of sprouts and for number with multiple sprouts (>1) at 2 weeks. One hundred %
- 20 signal solution induced multiple sprouts and dark incubation favoured sprouting as compared with the light regime after 1 wk of incubation. The exact cause of dark incubation favouring 100% signal solution is not understood. One hundred % signal solution was more effective than diluted signal solution when numbers of sprouted microtubers were counted after 1 wk. After 2 wk of incubation all treatments were
- 25 equally effective in causing sprouting but the signal solution and GA₃ solutions were most promotive of multiple sprouting which did not occur in the water soaking treatment and only in the control treatment incubated in the dark.

TABLE 4

**Effects of signal solution (SS) on dormancy breaking
of potato microtubers as compared to the known
dormancy breaker gibberellic acid (GA₃)**

Treatments	Number of sprouted microtubers		Number of multiple sprouted microtubers (2wk)	Mean number of sprouts ± SE
	1wk	2wk		
GA ₃ 500 mg l ⁻¹ + light	1/5	5/5	2/5	2.5 ± 0.5
GA ₃ 500 mg l ⁻¹ - light	5/5	5/5	4/5	2.5 ± 0.3
100 % SS + light	0/5	5/5	2/5	2.5 ± 0.5
100 % SS - light	5/5	5/5	3/5	2.3 ± 0.3
20 % SS + light	0/5	4/5	0	0
20 % SS - light	2/5	5/5	2/5	2.0 ± 0
12 % SS + light	0/5	4/5	1/4	2.0 ± 0
12 % SS- light	2/5	5/5	2/5	2.0 ± 0
6 % SS + light	0/5	4/5	2/4	2.0 ± 0
6 % SS - light	1/5	5/5	2/5	2.0 ± 0
Water + light	0/5	4/5	0	0
Water - light	2/5	5/5	0	0
Control + light	0/5	5/5	0	0
Control - light	1/5	5/5	1/5	2.0 ± 0

Tuber sprouting of potatoes is somewhat comparable to seed germination in the sense that the plant meristems are activated and the plant is beginning to grow, following quiescence. A signal molecule involved in bacteria-legume signalling was shown to be effective in breaking the dormancy of a plant (potato) that is distantly related to the legumes. LCOs therefore seem to have a broad effect on breaking the dormancy or quiescence of plants.

EXAMPLE 7

Effects of combinations of GA_3 and 100% signal solution in breaking dormancy of potato tubers

The data presented in Example 6 suggested that the 100% signal solution (and thus the LCO purified from *B. japonicum*) was effective in breaking microtuber dormancy. However, the microtubers used in the trial had been cold-stored for 8 wk. In this trial, the effects of signal solution were evaluated in combination with GA_3 on minitubers with only 3 wk of cold storage. It was also investigated whether the effect of 100% signal solution might be synergistic if used with GA_3 (500 mg l^{-1}).

Minitubers (20-30 g) with 3 wk cold storage treatment were soaked for 24 h in 500 mg l^{-1} GA_3 , 100% signal solution, or a mixture of the two. Another treatment involved successive soaking for 12 h each, in first GA_3 , and then signal solution. A control treatment without soaking was also performed. Eight minitubers were used per treatment which were applied at room temperature (20°C). Microtubers were observed after 2 wk and the number of sprouted minitubers and the number with multiple sprouts were counted.

All treatments were able to break minituber dormancy except the control (Table 5). The 500 mg l^{-1} GA_3 treatment alone or together with 100% signal solution, for 24 h, caused 100% sprouting, and significantly more multiple sprout formation than the other treatments. The 100% signal solution alone or in combination with 500 mg l^{-1} GA_3 were as effective as 500 mg l^{-1} GA_3 alone for breaking dormancy within 2 wk. However, less multiple sprouting occurred with 100% signal solution alone, or following 12 h GA_3 treatment, compared with the

GA₃ treatment alone or the combined GA₃ and Signal treatments. When working under the conditions tested, there were no clear synergistic effects of 100% signal solution in combinations with 500 mg l⁻¹ GA₃ on the number of sprouted tubers or multiple sprouts.

5

TABLE 5

Individual and combined effects of GA₃ and 100% signal solution (SS) on dormancy breaking of minitubers that had been cold-stored for 3 weeks

Treatments	Number of sprouted minituber	Number of minituber with multiple sprouts	Mean number of sprouts \pm SE
GA ₃ 500 mg l ⁻¹ 24 h	8/8	6/8	3.37 \pm 0.62
100% SS 24 h	7/8	2/7	1.33 \pm 0.42
GA ₃ 500 mg l ⁻¹ + 100% SS 12 h + 12 h	8/8	6/8	2.37 \pm 0.41
GA ₃ 500 mg l ⁻¹ + 100% SS Combination 24 h	8/8	6/8	3.75 \pm 0.67
Control	0/8	0/8	0/8

These results suggest that bacteria-legume signal molecules are effective in breaking the dormancy of potatoes. Taken together with the results presented above (i.e. Example 4 and Table 1) showing the effects of a pure LCO in breaking the quiescence of seeds and promoting growth of a variety of distantly

5

related plants, strongly supports the contention that LCOs are effective at breaking the dormancy of potatoes and promoting the activity of plant meristems in general.

EXAMPLE 8

5 **Effectiveness of GA₃ and signal solution, as compared to bromoethane and mechanical injuries, in breaking dormancy**

10 Bromoethane (BE) was reported to break potato tuber dormancy when applied as a fumigant and it was found that BE at a concentration of 0.2 ml l⁻¹ was the most effective (Coleman, 1983). Conventionally, large potato tubers are cut into small pieces, each containing an eye, to be used as seed pieces. To obtain quick and uniform sprout emergence, potato tubers should be cut at least 2 wk before planting (Slomnicki and Rylski, 1964). Mechanical injury is also shown to contribute to sprout induction. The objective of this experiment was to compare the effects of known dormancy-breaking treatments on microtubers and minitubers 1) BE; 2) GA₃; and 3) mechanical injury; with the newly identified dormancy breaker: LCOs.

20 Microtubers (200-500 mg) cold-stored for 8 wk and minitubers (20-35 g) cold-stored for 0, 2, or 8 wk were used for these experiments. BE (0.2 ml l⁻¹) and mechanical injury (cutting in half, microwaving at full power for 10 sec) were compared with GA₃ (500 mg l⁻¹), 100% signal solution, water soaking, and control treatments. Six microtubers or minitubers were used per treatment. Observations were made at 1, 2, 3, and 4 wk intervals and the number of sprouted tubers were counted. The evaluation period was extended because tubers with little or no cold storage treatment took longer to sprout.

30 GA₃ was the only agent which was able to break dormancy of minitubers that had not been cold-stored; 0/6 at 2 wk but 4/6 by 4 wk (minituber, 0 wk storage; Table 6). Minitubers with 2 wk cold storage that were treated with GA₃ also broke dormancy; 0/6 at 2 wk but 5/6 by 4 wk with 2/6 showing multiple shoots. Signal solution treatment of minitubers cold-stored for 2 wk caused 1/6 (with multiple shoots) minitubers to break dormancy after 4 wk. For minitubers cold-stored for 8 wk, all treatments (including water soaking and control), except the

microwaving, showed some sprouting the first week and multiple sprouting was evident in the GA₃ (6/6), 100% signal solution (4/6), and BE (1/6) treatments by 2 wk. Cutting caused sprouting in 9/12 cut halves by 2 wk but only a few multiple shoots (2/12) were evident by 3 wk.

097430-0520

TABLE 6

Sprouting and (multiple sprouting) performance on minitubers and microtubers cold-stored for 0, 2, 4, or 8 weeks after exposure to Bromoethane (0.2 ml l⁻¹), mechanical injury (cutting in half or microwaving), GA₃ (500 mg l⁻¹), 100% signal solution, water soaking, and control treatments.

Treatments	Minituber (CS8W)			Minituber (CS2W)			Minituber (NCS)			Microtuber (CS8W)						
	1wk	2wk	3wk	4wk	1wk	2wk	3wk	4wk	1wk	2wk	3wk	4wk				
GA ₃ 500mg l ⁻¹	5/6	6/6 (6)	6/6 (6)	6/6 (6)	0	0	3/6 (2)	5/6 (2)	0	0	3/6	4/6	4/6	5/6 (3)	6/6 (3)	6/6 (6)
100% Signal solution	4/6	6/6 (4)	6/6 (4)	6/6 (4)	0	0	0	1/6 (1)	0	0	0	0	1/6	2/6	4/6 (1)	6/6 (1)
Bromoethane 0.2 ml l ⁻¹	3/6	4/6 (1)	6/6 (4)	6/6 (4)	0	0	0	2/6	0	0	0	0	2/6	4/6	4/6	6/6
Cutting into halves	7/12	9/12	12/12 (2)	12/1 2 (2)	0	0	0	1/12	0	0	0	0	5/12	8/12	9/12	10/12
Microwaving	0	0	1/6	2/6	0	0	0	0	0	0	0	0	0	0	0	1/6
Water	4/6	5/6	6/6 (1)	6/6 (1)	0	0	0	0	0	0	0	0	2/6	4/6	6/6	6/6
Control	2/6	4/6	6/6	6/6	0	0	0	0	0	0	0	0	3/6	4/6	6/6	6/6

'CS' stands for cold storage; 'W' for week; N for no
'(')' denotes number of tubers with multiple sprouts

With microtubers cold-stored for 8 wk, dormancy-breaking occurred within the first week in all treatments except microwaving. After 2 wk, sprouting had progressed in all treatments; water (4/6) was similar to GA₃ solution (5/6), BE (4/6), signal solution (4/6), and cutting (8/12). Only GA₃ solution (3/6 at 2 wk, 6/6 at 4 wk) and signal solution (1/6 at 4 wk) caused multiple sprout formation.

Signal solution (100%) was effective in causing sprouting in minitubers with 2 or 8 wk cold storage and microtubers with 8 wk cold storage but it was ineffective on minitubers that had not been cold-stored. GA₃ and 100 % signal solution induced multiple sprouts from different eye-points unlike the control BE, or cutting treatments, that induced single sprouts only from the rose end. BE worked well in inducing single sprouts in minitubers and microtubers with 8 wk cold-storage but was not as effective as GA₃ for minitubers that had not been cold-stored.

Cutting minitubers or microtubers in half after 8 wk cold storage induced single sprouts on each cut half. This occurred quite efficiently in minitubers (9/12 in 2 wk, 12/12 in 4 wk) and somewhat less efficiently with microtubers (8/12 in 2 wk, 10/12 in 4 wk). Cutting was not effective in breaking dormancy in minituber without cold storage and worked poorly in minitubers that had been cold stored for only 2 wk (0/12 after 2 wk, 1/12 after 4wk). Sprouting from two halves was good, in the sense that by cutting minitubers or microtubers in half, two propagules, each with one sprout were derived although a very insignificant number of minituber halves (2/12) showed multiple sprouting. However, cutting was risky in that this sometimes provided opportunities for fungal or bacterial infection. Microwaving induced limited sprouting but only in minitubers or microtubers that had been cold-stored for 8 wk, and not in the minitubers with 0 or 2 wk cold storage. Microwaving caused some tuber damage that may account for the reduced sprouting observed.

In short, signal solution was effective in promoting both sprouting and multiple sprouting of mini- and micro-tubers and, of the tested treatments, only GA₃ was better.

EXAMPLE 9**Effectiveness of anti-ABA compared with other dormancy breaking treatments**

Anti-Abscisic acid (anti-ABA), the acetylenic analog of ABA, has never been used to induce sprouting in dormant potato tubers since it was first shown to be an ABA antagonist (Wilen *et al.*, 1993). However, anti-ABA has been used to terminate dormancy in canola seeds (PBI Bulletin, 1995). The objective of this experiment was to test anti-ABA for breaking dormancy in potato minitubers and compare its efficacy with other dormancy-breaking treatments.

Microtubers (200-600 mg) were cold-stored for 3 wk prior to the experiment. Seven microtubers were used per treatment. Treatments included 24 h soaks in anti-ABA or GA_3 (500 and 250 mg l^{-1} , respectively) applied alone or in combination, GA_3 (500 mg l^{-1} in combination with 100% signal solution, and water. Bromoethane (0.2 ml l^{-1}) and control treatments were also performed. Observations were made after 2 wk in the dark at room temperature (20°C). Data included number of sprouted microtubers and number of multiple sprouts. Means of sprout number were calculated only from microtubers that had sprouted

Anti-ABA alone and in successive treatments or in combination with GA_3 was effective in breaking microtuber dormancy (Table 7). Among the different treatments using anti-ABA and GA_3 the greatest mean number of sprouts (1.8 ± 0.48) occurred when microtubers were soaked in a mixed solution of 500 mg l^{-1} GA_3 and 500 mg l^{-1} anti-ABA for 24 h but it was not significantly different from the 500 mg l^{-1} GA_3 treatment (1.71 ± 0.28). The combined signal solution and GA_3 was not more effective than GA_3 alone and was less effective than any GA_3 and anti-ABA treatment in breaking dormancy.

TABLE 7

**Sprouting performance on microtubers cold-stored
for 3 weeks and evaluated at 2 weeks after
exposure to dormancy-breaking agents.**

Treatments	No. sprouted microtubers	Mean No. sprouts \pm SE
GA ₃ 500 mg l ⁻¹ 24 h	7/7	1.71 \pm 0.28
GA ₃ 500 mg l ⁻¹ + 100% SS 24 h	3/7	1.33 \pm 0.29
Anti-ABA 500 mg l ⁻¹ 24 h	5/7	1.4 \pm 0.24
Anti-ABA 250 mg l ⁻¹ 24 h	5/7	1.2 \pm 0.20
GA ₃ 500 mg l ⁻¹ 12 h + Anti-ABA 500 mg l ⁻¹ 12 h	6/7	1.33 \pm 0.21
GA ₃ 250 mg l ⁻¹ 12 h + Anti-ABA 250 mg l ⁻¹ 12 h	7/7	1.57 \pm 0.29
GA ₃ 500 mg l ⁻¹ + Anti-ABA 500 mg l ⁻¹ Combination 24 h	5/7	1.8 \pm 0.48
GA ₃ 250 mg l ⁻¹ + Anti-ABA 250 mg l ⁻¹ Combination 24 h	5/7	1.6 \pm 0.24
Bromoethane	1/7	1.0 \pm 0
Water	1/7	1.0 \pm 0
Control (no treatment)	0/7	0

The overall results with anti-ABA underline its importance as a potential dormancy-releasing agent, as much so as GA_3 . Anti-ABA and GA_3 both induced multiple sprouts but sprouts were longer after GA_3 than anti-ABA treatment. Both agents caused sprouts to emerge at various eyes over the tuber surface. However, the GA_3 -induced multiple sprouts were profusely branched; a group of sprouts protruded from each eye, while the anti-ABA-induced sprouts were singles. The mechanism of dormancy breaking by anti-ABA and GA_3 therefore was similar, but GA_3 appeared stronger. These agents should be tested on an equimolar basis in the future.

EXAMPLE 10

Harvests from minitubers sprouted using a range of dormancy-breaking treatments

There is only limited information on the relative yield performance of potato tubers that were treated with dormancy-breaking agents (Choudhury and Ghose, 1960; Slomnicki and Rylski, 1964). Yields from potato tubers that were treated with GA_3 at 25-100 mg l^{-1} (Choudhuri and Ghose, 1960) or 5-40 mg l^{-1} (Slomnicki and Rylski, 1964) were reduced compared with untreated controls. The objective of this experiment was to evaluate the effect of dormancy-breaking agents on subsequent yield in greenhouse pot trials.

Minitubers (20-35 g) that were cold-stored for 8 wk were given dormancy-releasing treatments including 24 h soaking in GA_3 (500 mg l^{-1}), 100% signal solution, or water. Other treatments included BE (0.2m l^{-1}), cutting in half, and the control. All minitubers were observed at 3 wk following treatment and the number of sprouts were noted at the time of planting. Five minitubers per treatment were individually planted into 11 x 12 cm plastic pots in the greenhouse. The potting mixture was 2:1 peat:perlite without fertilizer added. The pots were arranged in a complete randomized design and watered equally every alternate day. Harvest occurred after 60 d and tuber yields (number and fresh weight) were recorded.

GA₃ caused significantly more sprouts per minituber (4.2 ± 0.37) than the other treatments, with 100% signal solution (2.0 ± 0.31) and BE (1.8 ± 0.2) giving intermediate values, and water-soaking and cutting similar to the control (Table 8). The average number of tubers per plant was greatest in the GA₃ treatment (3.6); almost double that of other treatments that were not different from the control. Surprisingly, however, the mean fresh weight of tubers (per replicate i.e. pot basis) harvested from minitubers exposed to the 100% signal solution treatment was the greatest (34.97 g); greater than the control fresh weight and three times more than the GA₃ treatment. The size and shape of tubers harvested from the GA₃ treatment were small and more elongated than that of the control and other treatments. Yields from BE treated minitubers were significantly lower compared with controls. The cut halves each yielded almost the same as uncut controls and had similar fresh weight to control (28.41 vs 27.16). Two cut halves of each minituber together would effectively double control yield and bring the mean number of tubers into the GA₃ treatment range. However, cutting into halves posed a problem of infection and decomposition at the cut surfaces.

Thus, although signal solution is not as efficacious as GA₃ in breaking dormancy (as evaluated by the number of sprouts), it however is significantly more efficient than GA₃ in increasing the tuber yield. LCOs therefore appear as the best agents to promote dormancy breaking and yield increases in potato.

TABLE 8

Harvests after 60 d from minitubers that were forced to break dormancy by different methods.

Treatments	Mean number sprouts at planting \pm SE	Mean number of tubers produced per replicate	Mean fresh weight (g)
GA ₃	4.2 \pm 0.37	3.6*	11.13 d
Cutting into halves (1/2 minituber)	1.2 \pm 0.20	2.0	28.41 b
Bromoethane	1.8 \pm 0.20	2.0	14.01d
100% Signal soln.	2.0 \pm 0.31	1.8	34.97 a
Water	1.2 \pm 0.20	1.0	17.41 c
Control	1.0 \pm 0	2.0	27.16 b

Numbers represented by the same letter are not significantly different at the 0.05 level.

It shall be recognized therefore that agricultural compositions comprising at least one LCO and gibberellic acid (GA₃ and others known in the art) could be advantageously used in accordance with the methods of the present invention to break dormancy and/or quiescence of crop plants.

EXAMPLE 11

Other LCOs

Following the methods described above, the LCO most abundantly produced by *R. meliloti* (Nod Rm-V(C₁₆2, S)) was isolated and tested

on alfalfa (*Medicago sativa*) seeds. Briefly, 10 seeds were placed in a disk of filter paper on a petri plate. The filter paper was wetted with 5 ml of the appropriate LCO solution. Data were taken at 12 hour intervals upon the radicle (an embryonic root). The number of seed with an emerged radicle were counted. Each treatment was repeated four times. The data presented in table 9 indicate a clear acceleration of growth. In this case no standard for HPLC calibration was available, so a relative dilution series was used. In addition, a cluster of peaks specifically induced by the specific flavone of *Rhizobium leguminosarum* (bv *phaseoli* [strain 127K105]) were collected and tested on corn (*Zea mays*), red clover (*Trifolium repens*, Fabaceae) and pea (*Pisum sativum*, Fabaceae) (Table 10). In each case, a stimulation of seed germination was observed. Of note, *Rhizobium leguminosarum* produces a large number of LCOs. A subset of these LCOs was selected from a range of the HPLC profile where the LCOs from *B. japonicum* and *R. meliloti* did not occur. Taken together, these results clearly demonstrate that the promoting effects of LCOs on plant growth disclosed herein are observable with LCOs from different bacterial strains involved in bacteria-legume signalling. Consequently, the presented data strongly suggests that LCOs in general should demonstrate the same effects on seed germination, seedling emergence, growth, dormancy breakage and the like.

TABLE 9

**Effect of LCO isolated from *Rhizobium meliloti* (RCR 2011)
on germination of alfalfa after 24 h of treatment**

Treatment	Percent Germination
Control	16.7b
10 ⁻¹ dilution	26.6ab
10 ⁻² dilution	26.6ab
10 ⁻³ dilution	36.6a
LSD ($p < 0.05$)	19.2

In column numbers followed by same letter do not differ significantly by an ANOVA protected LSD test at $p < 0.05$

TABLE 10

Effect of LCOs of *Rhizobium leguminosarum* bv *phaseoli* (strain 127K105) on seed germination (%) of corn (after 48h), red clover (after 12h) and pea (after 48h) at 25°C

Treatment	Corn	Red Clover	Pea
Control	20a	43.3bc	26.6b
10 ⁻¹ dilution	26.6a	26.6c	26.6b
10 ⁻² dilution	60.0b	63.3ab	20.0b
10 ⁻³ dilution	20.0a	66.6a	73.3a
LSD ($p < 0.05$)	19.9	23.3	29.7

In column numbers followed by same letter do not differ significantly by an ANOVA protected LSD test at $p < 0.05$

EXAMPLE 12**Germination versus emergence**

Seeds of corn (cv Pioneer 3921) were surface sterilized in 2% sodium hypochlorite solution for 2 minutes and placed in 9 cm diameter Petri plates containing a sheet of filter paper soaked in 10 ml of the required test solution ($\text{LCO } 10^{-5} - 10^{-13}$). Water served as the control. Observations on germination, length of root primordia and shoot were taken after 72 h of incubation at 25°C. The data was analyzed for significance by an ANOVA protected LSD test using SAS system Version 6.1 (SAS Inc., Cary, NC, USA).

TABLE 11

Effect of lipo chitooligosaccharide [Bj Nod-V ($\text{C}_{18:1}$ MeFuc)] on germination of corn (*Zea mays* L.) after 72 h of treatment

Treatment	Percent germination	Length of root primordia (mm)	Length of shoot primordia (mm)
Control	46.6 a	32.3 a	4.6 a
$\text{LCO } 10^{-5}\text{M}$	80 bc	53.0 ab	12.3 ab
$\text{LCO } 10^{-7}\text{M}$	73.3 b	57.6 bc	15.0 ab
$\text{LCO } 10^{-9}\text{M}$	73.3 b	48.0 ab	9.6 a
$\text{LCO } 10^{-11}\text{M}$	100 c	78.6 c	21.0 b
$\text{LCO } 10^{-13}\text{M}$	80 bc	43.0 bc	8.3 a
LSD ($p < 0.05$)	22.6	24.3	11.0

In column numbers followed by same letter do not differ significantly by an ANOVA protected LSD test at $p < 0.05$.

Table 11 shows that incubation of corn seeds with LCO solution significantly improved the germination of corn and increased the length of both shoot and roots.

EXAMPLE 13**Seedling emergence-promoting effects of LCOs
under field conditions**

- Seeds of corn, cotton, beet, and soybean which showed
- 5 promising results under laboratory conditions were tested for seedling emergence under field condition. Seeds were surface sterilized with 2% sodium hypochlorite and soaked in different concentrations (10^{-5} , 10^{-7} , 10^{-9} M) of LCO solution for 12 h. Water served as the control. The study was conducted at the experimental field of the Macdonald campus of McGill University, Ste-Anne-de-
- 10 Bellevue, Quebec, Canada. The field was ploughed to a fine tilth. seeds were hand planted in 1 m rows at 2.5 to 3 cm deep with three replications per treatment. The percent seedling emergence was observed at six days after planting during which time at least 50% of the seeds emerged in the treatments. The data was analyzed for significance by an ANOVA protected LSD test using
- 15 SAS system Version 6.1 (SAS Inc., Cary, NC, USA).

TABLE 12

Effect of lipo chitooligosaccharide [Bj Nod-V ($C_{18,1}$ MeFuc)] on seedling emergence under field condition

Treatment	Corn	Cotton	Beet	Soybean
Control	41.6 a	6.6 a	26.6 a	16.6 a
LCO 10^{-5} M	80.0 b	16.6 a	28.3 a	26.6 ab
LCO 10^{-7} M	60.0 b	60.0 b	46.6 c	33.3 b
LCO 10^{-9} M	53.3 b	23.3 a	38.3 ab	63.3 c
LSD ($P < 0.05$)	35.7	28.0	16.3	16.6

In column numbers followed by same letter do not differ significantly by an ANOVA protected LSD test at $p < 0.05$

Table 12 shows that LCO [Bj Nod-V ($C_{18,1}$ MeFuc)] treatment enhanced the seedling emergence under field conditions of all the crop species studied. The best effect was observed in cotton where LCO at 10^{-7} M improved the emergence by more than 9 times as compared to the control. The effective concentration of LCO varied with the species.

Table 12 also validates the laboratory results presented herein by demonstrating that the stimulatory effects of LCOs are operating on four different crops under field conditions.

Thus the present invention provides agricultural compositions and methods by which LCO could be used to enhance the germination, seedling emergence, root growth and improve early growth of crops under laboratory or field conditions.

CONCLUSION

The present invention therefore provides evidence that, among other things: (1) lipo chitooligosaccharide (LCO) treatment enhances the seedling emergence of higher plant seeds (egs. *Z. mays*, *L. sativa*, *B. vulgaris*, *P. vulgaris*, *G. max*, *C. sativus*, *B. napus* and *M. sativa*); (2) lipo

chitooligosaccharide breaks the dormancy of potato (*Solanum tuberosum*)
minitubers and increases their yield; (3) lipo chitooligosaccharide improves
emergence and early growth, including root growth, of *Z. mays* giving a
competitive advantage over non treated ones; (4) lipo chitooligosaccharide
5 enhances the translocation of stored seed reserve; and (5) lipo
chitooligosaccharide enhances seed germination.

Although the present invention has been described
hereinabove by way of preferred embodiments thereof, it can be modified,
without departing from the spirit and nature of the subject invention as defined
10 in the appended claims.

REFERENCES

- Dakora et al., 1996, *Physiol & Molec Path* 49:1-20.
- De Jong et al., 1993, *Plant Cell* 5:615-620.
- 5 De Jong et al., 1997, *Phytopath* 87:588-593.
- Dénarié et al., 1996, *Annu Rev Biochem* 65:503-535.
- Durrant et al., 1991, *J. Plant Growth Reg.* 1:45-55.
- Fry et al., 1993, *Plant Phys* 103:1-5.
- Gillette et al., 1996, *J Bact* 178:2757-2766.
- 10 Heidstra et al., 1996, *New Phytol* 133:25-43.
- Kepczynski et al., 1992, *J. Plant Growth Reg.* 14:211-216.
- Long, 1996, *Plant Cell* 8:1885-1898.
- Long SR, 1989, *Cell* 56:203-214.
- Nap et al., 1990, *Science* 250:948-954.
- 15 Pavlova et al., 1992, *Plant Sci* 85:131-134.
- Peters et al., 1986, *Science* 233:977-980.
- Peters et al., 1990, *Mol. Plant-Microbe Interact.* 3:4-8.
- Röhrig et al., 1995, *Science* 269:841-843.
- Spaink HP, 1995, *Annu Rev Phytopath* 33:345-368.
- 20 Stacey et al., 1995, *Soil Biol Biochem* 27:473-483.
- Stacey G, 1995, *FEMS Microbiol Lett* 127:1-9.
- Stokkermans et al., 1995, *Plant Physiol* 108:1587-1595.
- Truchet et al., 1991, *Nature* 351:670-673.
- Verma DPS, 1992, *Plant Cell* 4:373-382.
- 25 Zhang et al., 1995, *Plant Physiol* 108:961-968.
- Zhang et al., 1997, *Soil Biol Biochem* 28:1579-1583.

EPO - DG 1

46

08. 09. 2000

WHAT IS CLAIMED IS:

(59)

1. Use of an agricultural composition for enhancing plant crop seed germination and/or seedling emergence and/or growth of a plant crop comprising a growth-promoting amount of at least one lipo
5 chitooligosaccharide (LCO) together with an agriculturally suitable carrier.
2. The use according to claim 1, wherein said plant crop is a non-legume.
3. The use according to claim 2, wherein said plant crop is selected from the group consisting of Poaceae, Cucurbitaceae, Malvaceae,
10 Asteraceae, Chenopodiaceae, Solanaceae and Brassicaceae.
4. The use according to claim 3, wherein said plant crop is selected from the group consisting of corn, cotton, cucumber, cantaloupe, lettuce, beet, canola and potato.
5. The use according to one of claims 1 to 4, wherein said
15 LCO is obtainable from a rhizobia selected from the group consisting of *Bradyrhizobium japonicum*, *Rhizobium meliloti* and *Rhizobium leguminosarum*.
6. The use according to claim 5, wherein said LCO is present in said composition at a concentration of between about 10^{-5} M to
20 about 10^{-14} M.
7. The use according to claim 5, wherein said LCO is present in said composition at a concentration of between about 10^{-6} M to about 10^{-12} M.
8. The use according to claim 5, wherein said LCO is
25 present in said composition at a concentration of between about 10^{-7} M to about 10^{-10} M.

9. The use according to one of claims 1 to 8, wherein said composition is effective in enhancing seed germination and/or seedling emergence and/or growth of a plant crop grown under field conditions.

5 10. The use according to claim 1, wherein said plant is a member of the Fabaceae family.

11. The use according to claim 10, wherein said plant is selected from the group consisting of soybean, bean, alfalfa and clover.

10 12. The use according to claim 10 or 11, wherein said LCO is obtainable from a rhizobia selected from the group consisting of *Bradyrhizobium japonicum*, *Rhizobium meliloti* and *Rhizobium leguminosarum*.

13. The use according to claim 12, wherein said LCO is present in said composition at a concentration of between about 10^{-6} M to about 10^{-14} M.

15 14. The use according to claim 12, wherein said LCO is present in said composition at a concentration of between about 10^{-6} M to about 10^{-12} M.

20 15. The use according to claim 12, wherein said LCO is present in said composition at a concentration of between about 10^{-7} M to about 10^{-10} M.

16. The use according to one of claims 10 to 15, wherein said composition is effective in enhancing seed germination and/or seedling emergence and/or growth of a plant crop grown under field conditions.

25 17. A use of a composition for breaking the dormancy and/or quiescence of a plant comprising a growth-promoting amount of at least one lipo chitoooligosaccharide (LCO) together with an agriculturally suitable carrier.

18. The use according to claim 17, wherein said plant is a member of the family of Solonaceae.

19. The use according to claim 18, wherein said plant is a potato.

5 20. The use according to claim 19, wherein said growth-promoting activity of said composition enables an increase in yield.

21. The use according to claim 19, further comprising gibberellic acid.

22. A method for enhancing seed germination and/or seedling emergence and/or growth of a plant, comprising a treatment in the vicinity of one of a seed, root or plant with a composition comprising an agriculturally effective amount of a lipo chitooligosaccharide (LCO) in admixture with an agriculturally suitable carrier medium, wherein said effective amount enhances seed germination and/or seedling emergence and/or growth of said plant in comparison to an untreated plant.

23. The method of claim 22, wherein said composition comprises at least one LCO at a concentration of between about 10^{-5} M to about 10^{-14} M.

24. The method of claim 23, wherein said composition comprises at least one LCO at a concentration of between about 10^{-6} M to about 10^{-12} M.

25. The method of claim 24, wherein said composition comprises at least one LCO at a concentration of between about 10^{-7} M to about 10^{-10} M.

26. A method for enhancing seed germination and/or seedling emergence and/or growth of a plant crop comprising incubating a rhizobial strain which expresses a lipo chitooligosaccharide (LCO) in the

vicinity of one of a seed and/or root of said plant such that said LCO enhances seed germination and/or seedling emergence and/or growth of said plant crop.

27. The method of claim 26, wherein said plant crop is a
5 non-legume.

28. The method of claim 27, wherein said plant crop is selected from the group consisting of Poaceae, Cucurbitaceae, Malvaceae, Asteraceae, Chenopodiaceae, Solanaceae and Brassicaceae.

29. The method of claim 28, wherein said plant crop is
10 selected from the group consisting of corn, cotton, cucumber, cantaloupe, lettuce, beet, canola and potato.


30. The method of one of claims 26 to 29, wherein said rhizobia is selected from *Bradyrhizobium japonicum*, *Rhizobium meliloti* and *Rhizobium leguminosarum*.

31. The method of one of claims 26 to 30, wherein said LCO
15 enhances seed germination and/or seedling emergence and/or growth of said plant grown under field conditions.

32. The method of claim 26, wherein said plant crop is a legume in the Fabaceae family and wherein said LCO enhances seed
20 germination and/or seedling emergence and/or growth of said legume grown under field conditions.

33. Use of an agricultural composition for enhancing a non-legume plant crop seed germination and/or seedling emergence and/or
25 growth of a plant grown under field conditions, comprising a growth-promoting amount of at least one lipo chitooligosaccharide (LCO) together with an agriculturally suitable carrier.

34. Use of an agricultural composition for enhancing a leguminous plant crop seed germination and/or seedling emergence and/or growth of a plant grown under field conditions, comprising a growth-promoting amount of at least one lipo chitooligosaccharide (LCO) together with an agriculturally suitable carrier.



United States Patent Application

COMBINED DECLARATION AND POWER OF ATTORNEY

As a below named inventor I hereby declare that: my residence, post office address and citizenship are as stated below next to my name; that

I verily believe I am the original, first and sole inventor (if only one name is listed below) or a joint inventor (if plural inventors are named below) of the subject matter which is claimed and for which a patent is sought on the invention entitled: COMPOSITION FOR ACCELERATING SEED GERMINATION AND PLANT GROWTH

The specification of which

- a. ☐ is attached hereto
b. ☒ was filed on as application serial no. and was amended on (if applicable) (in the case of a PCT-filed application) described and claimed in international no. PCT/CA99/00666 filed July 21, 1999 and as amended on (if any), which I have reviewed and for which I solicit a United States patent.

I hereby state that I have reviewed and understand the contents of the above-identified specification, including the claims, as amended by any amendment referred to above.

I hereby claim foreign priority benefits under Title 35, United States Code, § 119/365 of any foreign application(s) for patent or inventor's certificate listed below and have also identified below any foreign application for patent or inventor's certificate having a filing date before that of the application on the basis of which priority is claimed:

- a. ☐ no such applications have been filed.
b. ☒ such applications have been filed as follows:

FOREIGN APPLICATION(S), IF ANY, CLAIMING PRIORITY UNDER 35 USC § 119			
COUNTRY	APPLICATION NUMBER	DATE OF FILING (day, month, year)	DATE OF ISSUE (day, month, year)
Canada	2,243,669	21 July 1998	
ALL FOREIGN APPLICATION(S), IF ANY, FILED BEFORE THE PRIORITY APPLICATION(S)			
COUNTRY	APPLICATION NUMBER	DATE OF FILING (day, month, year)	DATE OF ISSUE (day, month, year)

I hereby claim the benefit under Title 35, United States Code, § 120/365 of any United States and PCT international application(s) listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States application in the manner provided by the first paragraph of Title 35, United States Code, § 112, I acknowledge the duty to disclose material information as defined in Title 37, Code of Federal Regulations, § 1.56(a) which occurred between the filing date of the prior application and the national or PCT international filing date of this application.

U.S. APPLICATION NUMBER	DATE OF FILING (day, month, year)	STATUS (patented, pending, abandoned)

I hereby claim the benefit under Title 35, United States Code § 119(e) of any United States provisional application(s) listed below:

U.S. PROVISIONAL APPLICATION NUMBER	DATE OF FILING (Day, Month, Year)

I acknowledge the duty to disclose information that is material to the patentability of this application in accordance with Title 37, Code of Federal Regulations, § 1.56 (reprinted below):

§ 1.56 Duty to disclose information material to patentability.

(a) A patent by its very nature is affected with a public interest. The public interest is best served, and the most effective patent examination occurs when, at the time an application is being examined, the Office is aware of and evaluates the teachings of all information material to patentability. Each individual associated with the filing and prosecution of a patent application has a duty of candor and good faith in dealing with the Office, which includes a duty to disclose to the Office all information known to that individual to be material to patentability as defined in this section. The duty to disclose information exists with respect to each pending claim until the claim is canceled or withdrawn from consideration, or the application becomes abandoned. Information material to the patentability of a claim that is canceled or withdrawn from consideration need not be submitted if the information is not material to the patentability of any claim remaining under consideration in the application. There is no duty to submit information which is not material to the patentability of any existing claim. The duty to disclose all information known to be material to patentability is deemed to be satisfied if all information known to be material to patentability of any claim issued in a patent was cited by the Office or submitted to the Office in the manner prescribed by §§ 1.97(b)-(d) and 1.98. However, no patent will be granted on an application in connection with which fraud on the Office was practiced or attempted or the duty of disclosure was violated through bad faith or intentional misconduct. The Office encourages applicants to carefully examine:

- (1) prior art cited in search reports of a foreign patent office in a counterpart application, and
- (2) the closest information over which individuals associated with the filing or prosecution of a patent application believe any pending claim patentably defines, to make sure that any material information contained therein is disclosed to the Office.

(b) Under this section, information is material to patentability when it is not cumulative to information already of record or being made of record in the application, and

- (1) It establishes, by itself or in combination with other information, a prima facie case of unpatentability of a claim;
- (2) It refutes, or is inconsistent with, a position the applicant takes in:
 - (i) Opposing an argument of unpatentability relied on by the Office, or
 - (ii) Asserting an argument of patentability.

A prima facie case of unpatentability is established when the information compels a conclusion that a claim is unpatentable under the preponderance of evidence, burden-of-proof standard, giving each term in the claim its broadest reasonable construction consistent with the specification, and before any consideration is given to evidence which may be submitted in an attempt to establish a contrary conclusion of patentability.

(c) Individuals associated with the filing or prosecution of a patent application within the meaning of this section are:

- (1) Each inventor named in the application;
- (2) Each attorney or agent who prepares or prosecutes the application; and
- (3) Every other person who is substantively involved in the preparation or prosecution of the application and who is associated with the inventor, with the assignee or with anyone to whom there is an obligation to assign the application.

(d) Individuals other than the attorney, agent or inventor may comply with this section by disclosing information to the attorney, agent, or inventor.

(e) In any continuation-in-part application, the duty under this section includes the duty to disclose to the Office all information known to the person to be material to patentability, as defined in paragraph (b) of this section, which became available between the filing date of the prior application and the national or PCT international filing date of the continuation-in-part application.

I hereby appoint the following attorney(s) and/or patent agent(s) to prosecute this application and to transact all business in the Patent and Trademark Office connected herewith:

Albrecht, John W.	Reg. No. <u>40,481</u>	Leonard, Christopher J.	Reg. No. <u>41,940</u>
Ali, M. Jeffer	Reg. No. <u>46,359</u>	Liepa, Mara E.	Reg. No. <u>40,066</u>
Anderson, Gregg I.	Reg. No. <u>28,828</u>	Lindquist, Timothy A.	Reg. No. <u>40,701</u>
Batzli, Brian H.	Reg. No. <u>32,960</u>	Lycke, Lawrence E.	Reg. No. <u>38,540</u>
Beard, John L.	Reg. No. <u>27,612</u>	Mayfield, Denise L.	Reg. No. <u>33,732</u>
Berns, John M.	Reg. No. <u>43,496</u>	McDonald, Daniel W.	Reg. No. <u>32,044</u>
Black, Bruce E.	Reg. No. <u>41,622</u>	McIntyre, Jr., William F.	Reg. No. <u>44,921</u>
Branch, John W.	Reg. No. <u>41,633</u>	Mitchem, M. Todd	Reg. No. <u>40,731</u>
Bremer, Dennis C.	Reg. No. <u>40,528</u>	Mueller, Douglas P.	Reg. No. <u>30,300</u>
Bruess, Steven C.	Reg. No. <u>34,130</u>	Nichols, A. Shane	Reg. No. <u>43,836</u>
Byrnie, Linda M.	Reg. No. <u>32,404</u>	Parsons, Nancy J.	Reg. No. <u>40,364</u>
Canipbell, Keith	Reg. No. <u>46,597</u>	Pauly, Daniel M.	Reg. No. <u>40,123</u>
Carlson, Alan G.	Reg. No. <u>25,959</u>	Phillips, Bryan K.	Reg. No. <u>46,990</u>
Caspers, Philip P.	Reg. No. <u>33,227</u>	Phillips, John B.	Reg. No. <u>37,206</u>
Chiapetta, James R.	Reg. No. <u>39,634</u>	Prendergast, Paul	Reg. No. <u>46,068</u>
Clifford, John A.	Reg. No. <u>30,247</u>	Pytel, Melissa J.	Reg. No. <u>41,512</u>
Coldren, Richard J.	Reg. No. <u>44,084</u>	Qualey, Terry	Reg. No. <u>25,148</u>
Daignault, Ronald A.	Reg. No. <u>25,968</u>	Reich, John C.	Reg. No. <u>37,703</u>
Daley, Dennis R.	Reg. No. <u>34,994</u>	Reiland, Earl D.	Reg. No. <u>25,767</u>
Dalglish, Leslie E.	Reg. No. <u>40,579</u>	Roberts, Fred	Reg. No. <u>34,707</u>
Daulton, Julie R.	Reg. No. <u>36,414</u>	Samuels, Lisa A.	Reg. No. <u>43,080</u>
DeVries Smith, Katherine M.	Reg. No. <u>42,157</u>	Schmaltz, David G.	Reg. No. <u>39,828</u>
DiPietro, Mark J.	Reg. No. <u>28,707</u>	Schuman, Mark D.	Reg. No. <u>31,197</u>
Edell, Robert T.	Reg. No. <u>20,187</u>	Schumann, Michael D.	Reg. No. <u>30,422</u>
Epp Ryan, Sandra	Reg. No. <u>39,667</u>	Scully, Timothy B.	Reg. No. <u>42,137</u>
Glance, Robert J.	Reg. No. <u>40,620</u>	Sebald, Gregory A.	Reg. No. <u>33,280</u>
Goggin, Matthew J.	Reg. No. <u>44,125</u>	Skoog, Mark T.	Reg. No. <u>40,178</u>
Golla, Charles E.	Reg. No. <u>26,896</u>	Spellman, Steven J.	Reg. No. <u>45,124</u>
Gorman, Alan G.	Reg. No. <u>38,472</u>	Stoll-DeBell, Kirstin L.	Reg. No. <u>43,164</u>
Gould, John D.	Reg. No. <u>18,223</u>	Sumner, John P.	Reg. No. <u>29,114</u>
Gregson, Richard	Reg. No. <u>41,804</u>	Swenson, Erik G.	Reg. No. <u>45,147</u>
Gresens, John J.	Reg. No. <u>33,112</u>	Tellekson, David K.	Reg. No. <u>32,314</u>
Hamer, Samuel A.	Reg. No. <u>46,754</u>	Trembath, Jon R.	Reg. No. <u>38,344</u>
Hamm, Curtis B.	Reg. No. <u>29,165</u>	Tuchman, Ido	Reg. No. <u>45,924</u>
Harrison, Kevin C.	Reg. No. <u>46,759</u>	Tunheim, Marcia A	Reg. No. <u>42,189</u>
Hertzberg, Brett A.	Reg. No. <u>42,660</u>	Underhill, Albert L.	Reg. No. <u>27,403</u>
Hillson, Randall A.	Reg. No. <u>21,838</u>	Vandenburgh, J. Derek	Reg. No. <u>32,179</u>
Holzer, Jr., Richard J.	Reg. No. <u>42,668</u>	Wahl, John R.	Reg. No. <u>33,044</u>
Johnston, Scott W.	Reg. No. <u>39,721</u>	Weaver, Karrie G.	Reg. No. <u>43,245</u>
Kadievitch, Natalie D.	Reg. No. <u>34,196</u>	Welter, Paul A.	Reg. No. <u>20,890</u>
Karjeker, Shaukat	Reg. No. <u>34,049</u>	Whipps, Brian	Reg. No. <u>43,261</u>
Kettelberger, Denise	Reg. No. <u>33,924</u>	Whitaker, John E.	Reg. No. <u>42,222</u>
Keys, Jeramie J.	Reg. No. <u>42,724</u>	Wickhem, J. Scot	Reg. No. <u>41,376</u>
Knearl, Homer L.	Reg. No. <u>21,192</u>	Williams, Douglas J.	Reg. No. <u>27,054</u>
Kowalchuk, Alan W.	Reg. No. <u>31,535</u>	Withers, James D.	Reg. No. <u>40,376</u>
Kowalchuk, Katherine M.	Reg. No. <u>36,848</u>	Witt, Jonelle	Reg. No. <u>41,980</u>
Lacy, Paul E.	Reg. No. <u>38,946</u>	Wu, Tong	Reg. No. <u>43,367</u>
Larson, James A.	Reg. No. <u>40,443</u>	Xu, Min S.	Reg. No. <u>39,536</u>
Leon, Andrew J.	Reg. No. <u>46,869</u>	Zeuli, Anthony R.	Reg. No. <u>45,255</u>

I hereby authorize them to act and rely on instructions from and communicate directly with the person/assignee/attorney/firm/ organization who/which first sends/sent this case to them and by whom/which I hereby declare that I have consented after full disclosure to be represented unless/until I instruct Merchant & Gould P.C. to the contrary.

Please direct all correspondence in this case to Merchant & Gould P.C. at the address indicated below:

Merchant & Gould P.C.
P.O. Box 2903
Minneapolis, MN 55402-0903



I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

200	Full Name Of Inventor	Family Name SMITH	First Given Name Donald	Second Given Name
0	Residence & Citizenship	City Sainte-Anne-de-Bellevue	State or Foreign Country Quebec, Canada CAX	Country of Citizenship Canada
1	Mailing Address	Address 2111 Lakeshore Road, P.O. Box 86	City Sainte-Anne-de-Bellevue	State & Zip Code/Country Quebec, H9X 3V9 Canada
Signature of Inventor 201:			Date: April 26, 2001	
200	Full Name Of Inventor	Family Name BO	First Given Name Pan	Second Given Name
0	Residence & Citizenship	City Sainte-Anne-de-Bellevue	State or Foreign Country Quebec, Canada CAX	Country of Citizenship China
2	Mailing Address	Address 2111 Lakeshore Road, P.O. Box 86	City Sainte-Anne-de-Bellevue	State & Zip Code/Country Quebec H9X 3V9 Canada
Signature of Inventor 202:			Date: May 3 2001	
300	Full Name Of Inventor	Family Name DENG	First Given Name Yinghai	Second Given Name
0	Residence & Citizenship	City Montreal	State or Foreign Country Quebec, Canada CAX	Country of Citizenship China
3	Mailing Address	Address 3655 Drummond	City Montreal	State & Zip Code/Country Quebec H3G 1Y6 Canada
Signature of Inventor 203:			Date: April 27, 2001	
400	Full Name Of Inventor	Family Name MIGNER	First Given Name Pierre	Second Given Name
0	Residence & Citizenship	City Sainte-Anne-de-Bellevue	State or Foreign Country Quebec, Canada CAX	Country of Citizenship Canada
4	Mailing Address	Address 21048 Lakeshore Road	City Sainte-Anne-de-Bellevue	State & Zip Code/Country Quebec H9X 1S2 Canada
Signature of Inventor 204:			Date: 25 avril 2001	
500	Full Name Of Inventor	Family Name ZHANG	First Given Name Feng	Second Given Name
0	Residence & Citizenship	City Sainte-Anne-de-Bellevue	State or Foreign Country Quebec, Canada CAX	Country of Citizenship Canada
5	Mailing Address	Address 7 Glenaladale Terrace	City Sainte-Anne-de-Bellevue	State & Zip Code/Country Quebec H9X 3V9 Canada
Signature of Inventor 205:			Date: April 24, 2001	

720	Full Name Of Inventor	Family Name <u>PRITHIVIRAJ</u>	First Given Name <u>Balakrishnan</u>	Second Given Name
0	Residence & Citizenship	City Sainte-Anne-de-Bellevue	State or Foreign Country Quebec, Canada <u>CA</u>	Country of Citizenship India
6	Mailing Address	Address 2111 Lakeshore Road, P.O. Box 86	City Sainte-Anne-de-Bellevue	State & Zip Code/Country Quebec H9X 3V9 Canada
Signature of Inventor 206: <u>B Prithiviraj</u>				Date: <u>26 April 2001</u>
720	Full Name Of Inventor	Family Name <u>HABIB</u>	First Given Name <u>Ahsan</u>	Second Given Name
0	Residence & Citizenship	City Sainte-Anne-de-Bellevue	State or Foreign Country Quebec, Canada <u>CA</u>	Country of Citizenship Bangladesh
7	Mailing Address	Address 2111 Lakeshore Road, P.O. Box 86	City Sainte-Anne-de-Bellevue	State & Zip Code/Country Quebec H9X 3V9 Canada
Signature of Inventor 207: <u>Affabir</u>				Date: <u>30/04/2001</u>